



# GEMTEC

[www.gemtec.ca](http://www.gemtec.ca)

**Environmental Impact Statement  
Proposed Plan of Subdivision  
3160 Carp Road  
Ottawa, Ontario**



# GEMTEC

[www.gemtec.ca](http://www.gemtec.ca)

Submitted to:

TLC Holdings Ltd.  
1380 Howie Road  
Carp, Ontario  
K0A 1L0

**Environmental Impact Statement  
Proposed Plan of Subdivision  
3160 Carp Road  
Ottawa, Ontario**

March 13, 2025  
Project: 102151.001

## EXECUTIVE SUMMARY

GEMTEC Consulting Engineers and Scientists Limited (GEMTEC) was retained by TLC Holdings Ltd. to complete an Environmental Impact Statement (EIS) in support of a proposed plan of subdivision for the property located at 3160 Carp Road, Ottawa, Ontario. The proponent is seeking to develop approximately 23 ha for a subdivision with road access from Carp Road.

To complete this EIS a desktop review and a series of field investigations were completed to identify the presence or absence of natural heritage features and species at risk (SAR) on-site. Field investigations were completed in spring and summer 2023.

Following completion of the desktop review and site investigations the following natural heritage features were identified on-site or within the study area: fish habitat and significant wildlife habitat for special concern and rare wildlife habitat (barn swallow, eastern wood-pewee, and wood thrush). The following SAR and their habitat were identified as having a potential to occur on-site: Blanding's turtle, eastern small-foot myotis, little brown myotis, tri-colored bat. No butternut trees were observed on-site.

Potential impacts to the natural heritage features were primarily associated with the loss of vegetation on-site due to potential future development. Potential impacts to significant wildlife habitat and fish habitat are primarily associated with loss of habitat and cumulative impacts to water quality through increases in nutrient and sediment loading.

Potential impacts to natural heritage features on-site can be mitigated through implementation of the recommended 15 m setback from the on-site watercourse.

Given the proposed development and minimal impact potential to Blanding's turtle and their habitat, it is GEMTEC's opinion that standard avoidance and mitigation measures will be sufficient to mitigate impacts of the proposed project and no ministry consultation is required.

Additionally, to provide protection to potential SAR and their habitat on-site, reptile and amphibian exclusion fencing should be installed around individual lots prior to any development or site alteration within the lot, to prevent the immigration of SAR turtles and other wildlife into the construction area. Should any SAR be discovered throughout the course of any development on-site, operations should stop and the species at risk biologist with the local MECP district should be contacted immediately for further direction.

## TABLE OF CONTENTS

1.0	INTRODUCTION.....	1
1.1	Purpose .....	1
1.2	Objective.....	1
1.3	Physical Setting.....	2
1.3.1	Land Use Context.....	2
2.0	METHODOLOGY.....	3
2.1	Desktop Review.....	3
2.2	Field Investigations.....	3
2.2.1	Ecological Land Classification.....	4
2.2.2	Breeding Bird Surveys.....	4
2.2.3	Headwater Drainage Feature Assessment .....	4
2.3	Data Analysis.....	5
3.0	EXISTING ENVIRONMENT.....	6
3.1	Ecoregion .....	6
3.2	Landforms, Soils and Bedrock Geology .....	6
3.3	Study Area Land Use .....	6
3.4	Carp River Watershed/Subwatershed Study and Carp Road Corridor Community Design Plan .....	7
3.5	Surface Water, Groundwater and Fish Habitat.....	8
3.5.1	Headwater Drainage Feature Assessment .....	8
3.6	Vegetation Communities .....	9
3.7	Wildlife .....	10
4.0	NATURAL HERITAGE FEATURES.....	11
4.1	Significant Wetlands .....	11
4.2	Significant Woodlands .....	11
4.3	Significant Valleylands.....	12
4.4	Significant Areas of Natural and Scientific Interest.....	12
4.5	Significant Wildlife Habitat .....	12
4.5.1	Habitats of Seasonal Concentrations of Animals .....	13
4.5.2	Rare Vegetation Communities .....	13
4.5.3	Specialized Habitats for Wildlife .....	13
4.5.4	Habitats of Species of Conservation Concern.....	13
4.5.5	Animal Movement Corridors .....	15
4.6	Fish Habitat .....	15
4.7	Species at Risk.....	16

5.0	PROPOSED PROJECT .....	17
6.0	IMPACT ASSESSMENT .....	18
6.1	Significant Wildlife Habitat .....	18
6.1.1	Habitats for Special Concern and Rare Wildlife Species SWH .....	18
6.2	Fish Habitat .....	19
6.2.1	Headwater Drainage Features .....	20
6.3	Species at Risk.....	21
6.3.1	Blanding’s Turtle.....	21
6.3.2	Eastern Red Bat .....	23
6.3.3	Eastern Small-footed Myotis .....	24
6.3.4	Hoary Bat .....	24
6.3.5	Little Brown Myotis .....	25
6.3.6	Silver-Haired Bat .....	26
6.3.7	Tri-colored Bat.....	26
6.4	Cumulative Impacts .....	27
7.0	RECOMMENDED AVOIDANCE AND MITIGATION MEASURES .....	28
7.1	Fish Habitat .....	28
7.1.1	Headwater Drainage Features .....	29
7.2	Significant Wildlife Habitat .....	30
7.2.1	Habitats of Species of Conservation Concern - Barn Swallow.....	30
7.3	Species at Risk.....	30
7.3.1	Blanding’s Turtle.....	30
7.3.2	Eastern Red Bat, Eastern Small-footed Myotis, Hoary Bat, Little Brown Myotis, Silver-Haired Bat & Tri-colored Bat.....	32
7.4	Wildlife .....	32
7.5	Best Practice Measures for Mitigation of Cumulative Impacts.....	33
8.0	CONCLUSIONS.....	34
9.0	LIMITATION OF LIABILITY.....	35
10.0	REFERENCES.....	36

**LIST OF TABLES**

Table 2.1	Summary of Field Investigations .....	4
Table 3.1	Vegetation Communities On-site.....	10

## LIST OF FIGURES

Figure 1 – Temporal Changes in Land Use within Study Area .....	7
Figure 4.1 Flow Chart Providing Directions of Management Option's (TRCA/CVC, 2014) .....	16

## LIST OF APPENDICES

Appendix A	Report Figures
Appendix B	Site Photographs
Appendix C	Report Summary Tables
Appendix D	Headwater Drainage Feature Assessment (GEMTEC, 2024)
Appendix E	MNRF General Habitat Descriptions – Blanding's Turtle

## 1.0 INTRODUCTION

GEMTEC Consulting Engineers and Scientists Limited (GEMTEC) was retained by TLC Holdings Ltd. to complete an Environmental Impact Statement (EIS) for the property municipally addressed as 3160 Carp Road, in the City of Ottawa, Ontario (hereafter referred to as “the subject property”). The location of the subject property is illustrated on Figure A.1 in Appendix A.

### 1.1 Purpose

The proponent is seeking to develop the approximately 23 hectare (ha) property for a future rural commercial/industrial subdivision. Based on *Section 4.7 – Environmental Protection* of the City of Ottawa Official Plan (Ottawa, 2022) an EIS is required showing that the proposed development will not negatively impact any potential natural heritage features, which may be present within the study area. The study area is defined as the property boundary and the adjacent lands encompassing an area of 120 m beyond the property boundary. The subject property and the extents of the study area are illustrated on Figure A.2 in Appendix A.

### 1.2 Objective

The 2024 Provincial Planning Statement (MMAH, 2024) issued under Section 3 of the Planning Act states that “development and site alteration shall not be permitted in: habitats of species at risk, significant wetlands, significant woodlands and significant wildlife habitat unless it has been demonstrated that there will be no negative impacts on the natural features or their ecological functions.” Similarly, the 2020 Provincial Planning Statement dictates that ‘development and site alteration shall not be permitted in fish habitat except in accordance with provincial and federal requirements.’”

The objective of the work presented herein is twofold; 1) to identify and evaluate the significance of any natural heritage features, as defined in the Provincial Planning Statement (MMAH, 2024), on the subject property and within the broader study area and; 2) to assess the potential impacts from the proposed plan of subdivision on any natural heritage features identified and to recommend appropriate and defensible mitigation measures to ensure the long-term protection of any natural heritage features identified.

To meet these objectives, the EIS presented herein has been completed in accordance with the following provincial and municipal regulations, policies and guidelines:

- Provincial Planning Statement (MMAH, 2024);
- Endangered Species Act (Ontario, 2007);
- Conservation Authorities Act (Ontario, 1990);
- Natural Heritage Reference Manual (OMNR, 2010);
- City of Ottawa Official Plan (Ottawa, 2022b); and
- City of Ottawa EIS Guidelines (Ottawa, 2023).

### **1.3 Physical Setting**

The subject property is municipally addressed as 3160 Carp Road, in the City of Ottawa, Ontario. The subject property currently consists of a mosaic of deciduous woodlands, agricultural field, thicket and cultural meadow. The site is bound to the northeast by Part of Lot 12, Concession 2 and to the northwest by 3186 Carp Road. To the southwest the site is bound by Carp Road and the rear lot lines of neighbouring properties on part of Lot 12, Concession 2. To the southeast the site is bound by lot lines of neighbouring properties on Lot 11, Concession 2.

#### **1.3.1 Land Use Context**

The subject property is situated within a larger rural area. The existing land use designation from the City of Ottawa is Rural Industrial and Logistics. The City of Ottawa zoning by-law is rural commercial zone, subzone 9 (RC9).

## 2.0 METHODOLOGY

### 2.1 Desktop Review

A desktop information gathering exercise was completed to aid in the scoping of field investigations and to gather information relating to natural heritage features which may be present on the subject project or within 1 km of the subject property. An additional component of the desktop review was to assess the potential presence of SAR to occur on the subject property or within the study boundary based on a review of publicly accessible occurrence records and a review of SAR habitat requirements and range maps.

Information regarding the potential presence of natural heritage features and SAR within the vicinity of the site was obtained from the following sources:

- Make a Map: Natural Heritage Areas (OMNRF, 2014a)
- Land Information Ontario (OMNRF, 2011);
- City of Ottawa Official Plan (City of Ottawa, 2022b)
- Ontario Geological Survey (OGS, 2019);
- Fisheries and Oceans Canada SAR Maps (DFO, 2023);
- Natural Heritage Information Centre Biodiversity Explorer (OMNRF, 2013);
- Breeding Bird Atlas of Ontario (Cadman et al., 2007);
- Ontario Herpetofaunal Atlas (Oldham and Weller, 2000);
- Wildlife Values Area (OMNRF, 2020a);
- Wildlife Values Site (OMNRF, 2020b); and
- Ontario Reptile and Amphibian Atlas (Ontario Nature, 2019).

### 2.2 Field Investigations

Field investigations were undertaken to describe in general, the natural and physical setting of the subject property with a focus on natural heritage features and to identify any potential SAR or their habitat that may exist at the subject property.

Field investigations completed in support of this EIS are outlined in Table 2.1 below. Photographs of site features taken during field investigations are provided in Appendix B.

**Table 2.1 Summary of Field Investigations**

Date	Time	Weather	Purpose
April 27, 2023	12:45 – 16:00	12°C, ~75% cloud cover, Beaufort 1, no precipitation	Headwater Drainage Feature Assessment
May 17, 2023	11:00 – 13:00	6°C, ~25% cloud cover, Beaufort 3, no precipitation	Headwater Drainage Feature Assessment
May 29, 2023	08:00 – 09:45	17°C, ~0% cloud cover, Beaufort 5, no precipitation	Breeding Bird Survey, Ecological Land Classification
June 15, 2023	08:15 – 10:15	18°C, ~60% cloud cover, Beaufort 4, no precipitation	Breeding Bird Survey, Ecological Land Classification
June 29, 2023	06:30 – 07:45	18°C, ~0% cloud cover, Beaufort 3, no precipitation	Breeding Bird Survey
July 26, 2023	07:15 – 08:45	19°C, ~25% cloud cover, Beaufort 1, no precipitation	Headwater Drainage Feature Assessment

### 2.2.1 Ecological Land Classification

Vegetation communities on the subject property were delineated during the desktop review stage of this EIS using publicly available air photos and confirmed in the field on May 29 and June 15, 2023, following the Ecological Land Classification System for Southern Ontario (Lee et al., 2008). Vegetation communities were confirmed in the field by employing the random meander methodology while documenting dominant vegetation species within the various vegetation community forms.

### 2.2.2 Breeding Bird Surveys

Breeding bird surveys were conducted on three occasions at four point count locations. Breeding bird surveys followed protocols from the Canadian Breeding Bird Surveys (Downes and Collins, 2003) and the Ontario Breeding Bird Atlas (Cadman, et al. 2007). Surveys were conducted no earlier than 30 minutes before sunrise and were completed within 5 hours of sunrise, to encompass peak song bird activity. Breeding bird surveys consisted of 5 minutes of passive listening in which all birds heard or seen within the survey period were recorded, including species, sex and breeding behaviour, if possible. A list of all avian species identified on-site is provided in Table C.1 in Appendix C.

### 2.2.3 Headwater Drainage Feature Assessment

A headwater drainage feature assessment (HDFa) was conducted to aid in the assessment of potential fish habitat on-site. Field data collection of headwater drainage features on-site followed the protocol outlined in Section 4: Module 11, “Unconstrained Headwater Sampling” from the Ontario Stream Assessment Protocol (Stanfield, 2017).

### **2.3 Data Analysis**

An evaluation of the significance of natural heritage features, the sensitivity of identified flora and fauna and the potential impacts posed by the proposed development was undertaken through an analysis of desktop and field investigation data using the approaches and criteria outlined in the following documents:

- Natural Heritage Reference Manual (OMNR, 2010);
- Significant Wildlife Habitat Technical Guide (OMNR, 2000);
- Significant Wildlife Habitat Ecoregion Criterion Schedules (OMNRF, 2015a); and
- Significant Wildlife Habitat Mitigation Support Tool (OMNRF, 2014b).

## **3.0 EXISTING ENVIRONMENT**

### **3.1 Ecoregion**

The site is situated Ecoregion 6E-11 (Lake Simcoe-Rideau), which extends from Lake Huron in the west to the Ottawa River in the east. The climate of Ecoregion 6E is categorized as humid, high to moderate temperate ecoclimate with a mean annual temperature range between 4.9°C to 7.8°C with annual precipitation ranging between 759 mm to 1,087 mm (Crins et al., 2009).

The eastern portion of the Ecoregion, which the subject property is located, is underlain by glaciomarine deposits as a result of the brief post-glacial incursion of salt water from the Champlain Sea along the St. Lawrence Valley. This Ecoregion falls with Rowe's (1972) Great Lakes-St. Lawrence Forest Region, including its Huron-Ontario and Upper St. Lawrence sections, and a small part of the Middle Ottawa Forest section (Crins et al., 2009).

### **3.2 Landforms, Soils and Bedrock Geology**

The topography of the site is relatively flat with a topographical high of 119 metres above sea level (mASL) in the southeast corner of the property to a topographical low of 108 mASL in the northeast of the property.

Two topographical landforms, as mapped by Chapman and Putnam (1984) are described on the subject property, sand plains and clay plains, both of the Ottawa Valley Clay Plains physiographic region. The majority of the property is comprised of sand plains, excluding a section of clay plains which occurs along the northeastern property boundary.

The Ontario Geological Survey (OGS, 2019) identifies two surficial soil units on the subject property, coarse-textured glaciomarine deposits and fine-textured glaciomarine deposits. The primary surficial soil unit is the coarse-textured glaciomarine deposits occurring in the southwest portion of the property comprised of littoral deposits consisting of sand, gravel, minor silt and clay. The remainder of the property consists of fine-textured glaciomarine deposits comprised of silt and clay, minor sand and gravel being massive to well laminated occurring in the northeast of the property.

Bedrock on the site is composed of the Ottawa Group; Simcoe Group; Shadow Lake Formation comprised of limestone, dolostone, shale, arkose, sandstone.

### **3.3 Study Area Land Use**

Figure 1 below provides an illustration of the temporal changes in land use within the study area from 1976, 2002, 2014 and 2022, based on aerial imagery from GeoOttawa.

In 1976, the subject property and surrounding lands were primarily populated with agricultural fields and small single family rural-residential dwellings buildings. Most development in the area was centred along Carp Road. Most of Kanata's urban area was not yet developed.

By 2002, the surrounding lands were still primarily agricultural, however, an increase in the number of small single family rural-residential dwellings were developed. Residential and commercial development within the study area primarily occurred south of the subject property.

By 2014, intensification of housing within the area to the south had reached present day extents. Development of smaller subdivisions continued to the southeast of the subject property.

By 2022, the remaining surrounding lands are in present day configuration.



**Figure 1 – Temporal Changes in Land Use within Study Area**

### **3.4 Carp River Watershed/Subwatershed Study and Carp Road Corridor Community Design Plan**

The Carp River Subwatershed Study (CRSWS) (Robinson, 2004) was completed, in part, to ensure that planning of future development proceeds in an environmentally sound manner. Specifically, the SWS aims to achieve this objective through making recommendations relating to how water resources and sub-watershed features, including their ecological functions are protected. The Carp River watershed encompasses an area of approximately 30,600 ha surrounding the former municipalities of West Carleton, Kanata and Goulbourn. The Carp River Watershed/Subwatershed Study (CRSWS) identifies opportunities and constraints for improvement of the Carp River Watershed while providing a series of Best Management Practices (BMPs) that may be implemented in order to protect, enhance or restore the environment.

Surface water within the subject property as identified in the Carp River SWS, is described as a warmwater, tolerant system with no sensitive, threatened or endangered species (Robinson, 2004). The portion of the Carp River that is located within the greater study area of this report was documented to have a combination of altered and natural conditions. The altered conditions along the section of the Carp River were representative of a long history of disturbance and reflected direct cattle access, bank erosion, excessive nutrients/abundant aquatic plant growth, and/or no vegetated buffer (Robinson, 2004).

The desktop review has identified a watercourse within the study area, however the CRSWS has not classified it as a cold-water stream. According to the CRSWS, the watercourse falls into the tolerant warmwater fish community, which is distributed throughout the main river between the Village of Kinburn and Richardson Side Road, “Glen Cairn Creek” and the upper portions of Feedmill, Huntley and Corkery Creeks. As such, under the recommendations provided by the CRSWS, the watercourse should receive a 15 m setback and revegetating up to 50% of the total stream length with native wood, riparian vegetation.

The Carp Road Corridor Community Design Plan (CRCCDP) is a Council approved guide to the long-term growth and development of the Carp Road Corridor. The CRCCDP provides guidelines for the day-to-day decision-making on land use planning and sets out the community’s priorities for the future (Ottawa, 2004). The Carp Road Corridor extends from Stittsville to Fitzroy Harbour and is a significant rural employment area. Schedule 2 of the CRCCDP identifies portions of the subject property as a moderate recharge area.

### **3.5 Surface Water, Groundwater and Fish Habitat**

Surface water on the subject property consists of one watercourse, as well as various headwater drainage features (HDFs).

The on-site watercourse originates along the southeastern property boundary and continues along the agricultural field in a northeasterly direction for approximately 1.5 km before discharging into the Carp River.

As a component of the HDFA, fish habitat was assessed within the on-site HDFs. No fish species were observed and the HDFs were determined to provide limited fish habitat due to limited hydroperiod and limited in stream structure, vegetation or habitat.

Groundwater investigations were not completed in support of this EIS.

#### **3.5.1 Headwater Drainage Feature Assessment**

A headwater drainage feature assessment (HDFA) was conducted for all un-named ephemeral watercourses on-site. Based on the desktop review and the site surveys, six HDFs occur on-site and are identified as H1, H2, H3, H3A, H3B, and H4. All six HDFs are illustrated on Figure A.2 in Appendix A.

H1 originates on-site near the northwestern property boundary and travels on-site for approximately 50 m before continuing off-site in a north-eastern direction along the agricultural fields, eventually discharging into the Carp River. HDF1 is likely a remnant of historical agricultural drainage efforts.

H2 is an isolated feature that originates in the north of the property and flows in a southeast direction for approximately 50 m before abruptly ending. HDF2 does not have any connection with other surface water features upstream or downstream.

H3 originates in the eastern extent of the property and flows through the cultural thicket in a northwest direction for approximately 460 m before discharging into a vertical drain. Based on site observations, the vertical drain appeared to be allowing the water to pool and reinfiltrate below ground. It did not appear to be moving laterally across the site (i.e. like a tile drain). As such it is assumed that H3 (and H3A and H3B) do not have direct connectivity to any downstream surface water features. They are considered unconnected features,

H3A originates in cultural thicket in the northeast of the property and flows in a northern direction before discharging into the downstream extent of HDF3, and ultimately, the vertical drain.

H3B is located within the eastern-central portion of the property and originates as an off-shoot from HDF3, within the cultural thicket. HDF3B continues from the confluence for approximately 20 m in a northeast direction before abruptly ending before approaching the agricultural fields.

H4 originates in the southeastern corner of the property and flows in a southeastern direction for approximately 35 m before discharging into the permanent watercourse along the agricultural field.

The evaluation, classification, and management recommendations for each HDF, as derived from the Guidance Document (CVC/TRCA, 2014) are provided in the HDFA for the property in Appendix D.

### **3.6 Vegetation Communities**

Vegetation communities on-site were confirmed by GEMTEC in 2023, following protocols utilized in the Southern Ontario Ecological Land Classification System (Lee et al., 2008). Vegetation at the site represents a mosaic of deciduous forest/woodland, cultural meadow, open agriculture, and thicket communities. Table 3.1 below provides a summary of the various vegetation communities identified on-site while Figure A.3 in Appendix A provides an illustration of the various vegetation communities.

**Table 3.1 Vegetation Communities On-site**

ELC Type	Description	Size (ha)
Fresh – Moist Lowland Deciduous Forest (FODM7)	Located in eastern portion of the property is a deciduous forest, dominated by American elm ( <i>Ulmus americana</i> ) and Manitoba maple ( <i>Acer negundo</i> ), with other common constituents including trembling aspen ( <i>Populus tremuloides</i> ) and willow species ( <i>Salix</i> sp.). The shrub layer was heavily dominated by buckthorn ( <i>Rhamnus</i> sp.) and included honeysuckle ( <i>Lonicera periclymenum</i> ). The herbaceous layer consisted of grasses.	2.7
Mixed Meadow (MEM)	Located in the southwestern half of the property is a mixed meadow dominated by grasses. Other common constituents included yellow salsify ( <i>Tragopogon dubius</i> ), orange hawkweed ( <i>Pilosella aurantiaca</i> ), wild parsnip ( <i>Pastinaca sativa</i> ), clover ( <i>Trifolium</i> sp.), vetch ( <i>Vicia</i> sp.), common daisy ( <i>Bellis perennis</i> ), common milkweed ( <i>Asclepias syriaca</i> ), meadow hawkweed ( <i>Hieracium caespitosum</i> ), and bladder campion ( <i>Silene vulgaris</i> ). Few trees were scattered within this community, consisting of white spruce ( <i>Picea glauca</i> ) and Scots pine ( <i>Pinus sylvestris</i> ).	14.1
Buckthorn Deciduous Shrub Thicket (THDM2-6)	Located in the eastern portion of the property is a deciduous thicket dominated by buckthorn and red-osier dogwood ( <i>Cornus sericea</i> ). Other common constituents included Manitoba maple, willow, and honeysuckle. The herbaceous layer consisted of grasses.	1.4
Meadow (ME)	<p>Located in the north of the property is a meadow community. Vegetation was dominated by tall grasses, goldenrod (<i>Solidago</i> sp.), brambles (<i>Rubus</i> sp.), wild parsnip, virgin's bower (<i>Clematis virginiana</i>), and dog-strangling vine (<i>Vincetoxicum rossicum</i>).</p> <p>Within the eastern extent of this community is a small wetland inclusion, characterized by grasses and sedges, bulrush (<i>Scirpoides holoschoenus</i>), cattail (<i>Typha</i> sp.), red-osier dogwood, purple loosestrife (<i>Lythrum salicaria</i>), jewelweed (<i>Impatiens capensis</i>), bedstraw (<i>Galium</i> sp.), and bittersweet nightshade (<i>Solanum dulcamara</i>).</p> <p>The wetland inclusion is not considered as its own community as it does not meet the minimum size requirements of 0.5 ha, as per the guidelines set out in the Ecological Land Classification System (Lee et al., 2008).</p>	0.5
Open Agriculture (OAG)	Located in the east section of the property is an open agricultural field, comprised of corn.	4.4

### 3.7 Wildlife

Wildlife observed on-site and within the study area during field investigations completed in 2023 are summarized in Table C.1 in Appendix C.

## 4.0 NATURAL HERITAGE FEATURES

Natural heritage features are defined in the PPS as “features and areas, including *significant wetlands, significant coastal wetlands, fish habitat, significant woodlands south and east of the Canadian Shield, significant valleylands south and east of the Canadian shield, significant habitats of endangered species and threatened species, significant wildlife habitat and significant areas of natural and scientific interest*, which are important for their environmental and social values as a legacy of the natural landscape of an area”.

Natural heritage features identified in Section 4 below are illustrated on Figure A.4 in Appendix A.

### 4.1 Significant Wetlands

As described in the Natural Heritage Reference Manual (OMNR, 2010), wetlands mean “lands that are seasonally or permanently covered by shallow water, as well as lands where the water table is close to or at the surface.” While *significant* in regards to wetlands means “an area identified as provincially significant by the Ontario Ministry of Natural Resources and Forestry using evaluation procedures established by the Province, as amended from time to time.”

No PSW or local wetlands were identified on-site during the desktop review, nor were they identified on-site. As such, significant wetlands are not discussed or evaluated further in this EIS.

### 4.2 Significant Woodlands

Significant woodlands are defined in the natural heritage reference manual (OMNR, 2010) as “an area which is ecologically important in terms of features such as species composition, age of trees and stand history; functionally important due to its contribution to the broader landscape because of its location, size or due to the amount of forest cover in the planning area; or economically important due to site quality, species composition, or past management history.”

At the local scale, significant woodlands are defined and designated by the local planning authority. Generally, most planning authorities have defined significant woodlands as any woodland that contains any of the four criteria listed in Section 7.2 of the natural heritage reference manual (OMNR, 2010), including: woodland size, ecological functions, uncommon characteristics and economic and social functional values. Furthermore, the City of Ottawa provides a supplementary document *Significant Woodland: Guidelines for Identification, Evaluation, and Impact Assessment* (Ottawa, 2022d) to evaluate woodlands and ensure compliance with the city’s policies.

As outlined in *Significant Woodlands: Guidelines for Identification, Evaluation and Impact Assessment* (Ottawa, 2022b), rural area woodlands are to be identified and evaluated using all the natural heritage resource manual (OMNR, 2010) criteria. Table C.2 in Appendix C, presents the screening rationale for significant woodlands applied in this EIS. For comparison of woodland criteria used in Table C.2, it is assumed that the woodland coverage within the planning area (City

of Ottawa – Rural Planning Area – Ottawa West) is between 30% and 60% of the land area, therefore the minimum woodland size for determining significance is 50 ha or greater, based on the guidance outlined in the natural heritage reference manual (OMNR, 2010).

Based on the NHRM (OMNR, 2010) screening criteria presented in Table C.2 in Appendix C, significant woodlands are not present on-site. As such, significant woodlands are not present on-site or within the study area and are therefore not discussed or evaluated further in this EIS.

#### **4.3 Significant Valleylands**

Valleylands are defined in the natural heritage reference manual (OMNR, 2010) as ‘a natural area that occurs in a valley or other landform depression that has water flowing through or standing for some period of time’. The identification and evaluation of significant valleys lands in Ontario is based on the recommended criteria from the MNRF and is the responsibility of local planning authorities.

In Southern Ontario, conservation authorities have identified valleylands as part of their regulation mapping (i.e., floodplain mapping); however, where valleys lands have not been defined, their physical boundaries are generally determined as the ‘top-of-bank’ or ‘top-of-slope’ associated with a watercourse. For less well-defined valleys, the physical boundary may be defined by riparian vegetation, flooding hazard limits, ordinary high-water marks or the width of the stream meander belt (OMNR, 2010).

As discussed in Section 3.2, the site is relatively flat, furthermore no valleylands were identified on-site during the desktop review or the site investigations. As such significant valleylands are not discussed or evaluated further in this EIS.

#### **4.4 Significant Areas of Natural and Scientific Interest**

The MNRF identifies two types of areas of natural and scientific interest (ANSI) in Ontario: life sciences ANSIs typically represent significant segments of Ontario’s biodiversity and natural landscapes, while earth science ANSIs typically represent significant examples of bedrock, fossils or landforms in Ontario (OMNR, 2010).

No ANSI have been identified on-site or adjacent to the site during the desktop review or during site investigations. Therefore, ANSI are not discussed or evaluated further in this EIS.

#### **4.5 Significant Wildlife Habitat**

The natural heritage reference manual (OMNR, 2010), in combination with the significant wildlife habitat technical guide (MNRF, 2000) and the significant wildlife habitat ecoregion criterion schedules (MNRF, 2015) were used to identify and evaluated potential significant wildlife habitat on-site. The significant wildlife habitat is broadly categorized as habitats of seasonal concentration of animals, rare vegetation communities, specialized habitats for wildlife, habitats of species of

conservation concern and animal movement corridors. Table C.3, C.4, C.5 and C.6 in Appendix C, provide the screening rationale for each category of significant wildlife habitat, respectively.

#### **4.5.1 Habitats of Seasonal Concentrations of Animals**

Seasonal concentration areas are habitats where large numbers of species congregate at one particular time of the year. The significant wildlife habitat technical guides (OMNR, 2000) and significant wildlife habitat ecoregion criterion schedules (OMNRF, 2015a) identify 11 types of seasonal concentration habitats that may be considered significant wildlife habitat. These 11 types of seasonal habitat are presented in Table C.3 in Appendix C, including a brief description of the rationale as to why they are or are not assessed further in this EIS.

Following review of Table C.3 in Appendix C, no habitats of seasonal concentrations of animals were identified on-site.

#### **4.5.2 Rare Vegetation Communities**

Rare vegetation communities in the province are described generally as those with an S1 to S3 ranking by the NHIC, and typically include communities such as sand barrens, alvars, old growth forests, savannahs and tallgrass prairies.

The vegetation communities identified on-site and described in Section 3.4 of this report are not ranked by the NHIC as S1, S2 or S3 and are therefore not considered to be rare vegetation communities. As such, rare vegetation communities are not discussed or evaluated further in this EIS.

#### **4.5.3 Specialized Habitats for Wildlife**

Specialized wildlife habitats are microhabitats that provide a critical resource to some groups of wildlife. The significant wildlife habitat technical guide (OMNR, 2000), defines eight specialized habitats that may constitute significant wildlife habitat, these eight types of specialized wildlife habitats are evaluated in Table C.4 in Appendix C.

Following review of Table C.4 in Appendix C, no specialized habitat for wildlife has been identified on-site or within the study area.

#### **4.5.4 Habitats of Species of Conservation Concern**

Provincial rankings are used by the Natural Heritage Information Centre to set protection priorities for rare species, similar to those described in Section 4.5.2 above for vegetation communities. Provincial rankings (S-ranks), are not legal designations such as those used to define the various protection statuses of species at risk, they are only intended to consider factors within the political boundaries of Ontario that might influence a particular species abundance, distribution or population trend.

Based on the guidance provided in the Significant Wildlife Habitat Ecoregion Criterion Schedules (MNRF, 2015), when a plant or animal element occurrence is recorded for any species with an S-rank of S1 (extremely rare), S2 (very rare), S3 (rare to uncommon) or SH (historically present), the corresponding vegetation ecosite is considered to provide *candidate* habitat for species of conservation concern and further consideration within the EIS is warranted.

The Significant Wildlife Habitat Ecoregion Criterion Schedules (OMNRF, 2015), provides five general habitat types known to support a wide range of species of conservation concern in Ontario. The five general habitat types for Ecoregion 6E-11 are provided in Table C.5 in Appendix C, including a brief rationale as to why they are or are not considered further in this EIS. Following review of Table C.5 in Appendix C, one habitat of species of conservation concern has been identified on-site, habitat for special concern and rare wildlife species for barn swallow, eastern wood-pewee and wood thrush.

#### 4.5.4.1 Special Concern and Rare Wildlife Species SWH

Based on observation data from the field investigations, one species of special concern has been identified on-site or within the broader study area, barn swallow. Based on observation data from the NHIC database, two species of special concern have been identified on-site and within the broader study area, eastern wood-pewee and wood thrush. No other species of special concern or rare wildlife species were identified on-site or within the broader study area.

The barn swallow (*Hirundo rustica*) is medium-sized, insectivorous bird with an S-rank of S4B (uncommon but not rare, breeding population) and is listed as a species of special concern in Ontario. No historical observations for barn swallow have been identified by the NHIC database within the study area. However, barn swallow were observed on-site foraging over the meadow and agricultural field (ELC codes: MEM and OAG) during the breeding bird site investigation on June 5, 2023. Given the observed sighting and availability of foraging habitat on-site, there is a high potential for barn swallow to occur on-site. However no suitable nesting sites or structures occurs on-site.

The eastern wood-pewee is a small flycatcher bird with an S-rank of S4 (uncommon but not rare) in Ontario. The NHIC identified the eastern wood-pewee as having historically occurred within 1 km of the site; however, eastern wood-pewee were not observed on-site during the breeding bird surveys or other fields investigations. Eastern wood-pewee is a woodland species that is often found near clearings and edges. Given the mosaic of woodland and open habitat on-site and the eastern wood-pewee's affinity for clearings and edges, there is a moderate chance for the eastern wood-pewee or suitable habitat to occur on-site.

The wood thrush is a medium-sized songbird with an S-rank of S4 (uncommon but not rare) and is listed as a species of special concern in Ontario. The NHIC identified the wood thrush as having historically occurred within 1 km of the site; however, wood thrush were not observed on-site

during the breeding bird surveys or other field investigations. Wood thrush is a woodland species often found in moist, deciduous hardwood or mixed forests stands, with dense deciduous undergrowth and tall trees. Given the forested community on-site and within the study area, there is a moderate chance for the wood thrush or suitable habitat to occur on-site.

#### **4.5.5 Animal Movement Corridors**

Animal movement corridors are elongated areas used by wildlife to move from one habitat to another and allow for the seasonal migration of animals (OMNRF, 2015). The Significant Wildlife Habitat Ecoregion Criterion Schedules for Ecoregion 6E-11 (OMNRF, 2015), identifies two types of animal movement corridor: amphibian movement corridors and deer movement corridors. As per guidance presented in MNRF, 2015, animal movement corridors should only be identified as significant wildlife habitat when a *confirmed or candidate* significant wildlife habitat has been identified by the MNRF district office or by the regional planning authority.

Following review of Table C.6 in Appendix C, no animal movement corridors have been identified on-site. Furthermore, the MNRF has not identified any animal movement corridors on the publicly available data sets for wildlife values area (OMNRF, 2020a) or wildlife values site (OMNRF, 2020b). As such, animal movement corridors are not discussed or evaluated further in this EIS.

#### **4.6 Fish Habitat**

The protection of fish and fish habitat is a federal responsibility and is administered by the Department of Fisheries and Oceans Canada (DFO). Fish habitat as defined in the Fisheries Act (Canada, 1985) means, “spawning grounds and nursery, rearing food supply and migration areas on which fish depend directly or indirectly in order to carry out their life processes.”

When development is unable to avoid resulting in the harmful alteration, disturbance or destruction of fish habitat from typical project impacts such as temperature change, sedimentation, infilling, reduction of nutrient and food supply, etc., an authorization under the Fisheries Act is required for the project to proceed.


A fisheries assessment was not conducted as part of this EIS, until such a time that a fisheries assessment is completed, the unnamed watercourse is assumed to provide fish habitat for small-bodied fish species.

Given the limited hydrology of the HDFs on-site, (all were dry but the third visit), and the limited connectivity to downstream features, the HDFs on-site are not considered to provide direct fish habitat.

Impacts to fish habitat on-site are discussed in Section 6.

#### 4.6.1.1 Headwater Drainage Feature Assessment

An HDFA was conducted to assess the contribution of un-mapped and un-named watercourses on-site to fish habitat. The full HDFA report is provided in Appendix D. Six HDFs were identified on-site and are illustrated on Figure A.2 in Appendix A. The watercourses on-site were classified based on the information collected during site investigations pertaining to hydrology, riparian habitat, fish and fish habitat and terrestrial components. Using the linking classification management flow chart provided by the TCRA and CVA (2014), illustrated in Figure 4.1 below, the characteristics of the on-site watercourses were used to determine the management recommendations.



**Figure 4.2 Flow Chart Providing Directions of Management Option's (TRCA/CVC, 2014)**

#### 4.7 Species at Risk

The probability of occurrence for species at risk to occur on-site and within the broader study area was determined through the desktop review stage of this EIS, as described in Section 2.1, and through the site-specific surveys conducted as part of this EIS, outlined in Section 2.2.

Table C.7 in Appendix C, provides a summary of all species at risk which were determined to have the potential to occur on-site or within the broader study area, their protection status under the provincial Endangered Species Act (Ontario, 2007), their regional distribution, their probability of occurrence and a brief rationale of that probability. Impacts to endangered or threatened SAR determined to have a moderate or high potential to occur on-site or within the broader study area are discussed further in the Section 6.3.

## 5.0 PROPOSED PROJECT

The proposed project, assessed for potential impacts on the natural heritage features determined to be present within the broader study area, is a subdivision application for the development of the approximately 23 ha property.

The proposed development at the site will consist of 12 lots with sizes ranging from about 1.1 ha to 4.8 ha in size, and the creation of a road. One of the proposed lots (either Block 1, 11 or 12) is proposed to be severed prior to the subdivision application. Access to the proposed subdivision will be from Carp Road. The proposed plan of subdivision is provided on Figure A.5.

Stormwater management will be provided for quality and quantity and is proposed to occur on Block 7. Post-development, stormwater runoff will be captured by a system of ditches/swales and conveyed to the proposed SWM facility on Block 7 before discharging to the existing permanent watercourse located on-site.

The proposed SWM facility has been designed with adequate storage volume to detain flows exceeding pre-development levels for up to and including the 100-year design storm event. The outlet channel located on the northside of Block 7 will convey stormwater runoff from the municipal right-of-way to the proposed SWM facility. The outlet channel will be designed in keeping with the recommendations for an enhanced grass swale and will provide pretreatment and attenuation for runoff prior to discharging into the SWM facility. The implementation of vegetated roadside ditches and rear yard ditches/swales, a flat bottom outlet channel, a dry pond, on-site block level quality controls, will provide a reasonable amount of quality control for stormwater runoff generated from the municipal right-of-way and future development blocks.

Future components of the proposed project considered in the impact assessment presented in Section 6 include: tree clearing and vegetation grubbing, fill placement and elevation grading, road construction, laneway construction, drilling of a groundwater well and septic system installation, excavation and pouring of a foundation, construction of commercial and light industrial buildings and general landscaping activities.

The timeline for the proposed project, from lot creation to completion of construction is currently unknown. For the purpose of assessing impacts to natural heritage features, it is assumed in this EIS that the creation of individual lots will happen in the near-term and will not result in any physical alterations to the natural environment of the site and the broader study area. Future construction of each lot is assumed to occur over a several year period, and that the construction of any lot will be completed such that the duration of any potential impacts on the natural environment during construction will be approximately six to nine months.

## 6.0 IMPACT ASSESSMENT

Potential impacts to natural heritage features on-site and within the broader study area are assessed for direct, indirect and cumulative effects based on the proposed project outlined in Section 5. Natural heritage features identified in Section 4 of this report as present or likely to be present are discussed in the subsections below.

Potential effects to the environment of the site from the proposed development outlined in Section 5 include: vegetation removal, habitat fragmentation and loss, disturbance of the natural soil mantle, increased noise generation, increased human disturbance, increase storm water generation, increased nutrient loading to adjacent surface water features, increase in impervious surface and short-term increases in sedimentation and/or erosion.

### 6.1 Significant Wildlife Habitat

The potential presence of *candidate* significant wildlife habitat on-site and within the study area was evaluated in Section 4.5. As a result of this assessment one significant wildlife habitats were determined to be present on-site or within the study area; *candidate* special concern and rare wildlife species SWH for barn swallow, eastern wood-pewee and wood thrush.

Potential impacts to each type of SWH are discussed in greater detail in the following subsections, while mitigation measures intended to prevent such impacts are presented in Section 7.

#### 6.1.1 Habitats for Special Concern and Rare Wildlife Species SWH

##### 6.1.1.1 Barn Swallow

Barn swallow (*Hirundo rustica*) is medium-sized, insectivorous bird listed as a species of special concern in Ontario.

Barn swallows typically build their nests out of mud on ledges or walls on barns or other human made structures. Natural sites, including cliffs and caves are rarely used for nesting (Cadman et al., 2007). Foraging occurs within fields and ponds. Barn swallows are less common in highly urban area and areas with higher forest cover (Cadman et al., 2007).

No historical occurrences have been reported for barn swallow within 1 km of the subject property; however, they were observed foraging during the site investigations.

Impacts to barn swallow and their habitat from the proposed development is limited to the open agricultural field and meadow habitats (ELC codes OAG, MEM, and ME), which may provide suitable foraging habitat. No nests or suitable nesting structures were directly observed on-site. Suitable nesting structures may occur off-site within the study area. Additionally, an abundance of habitat for foraging barn swallow is present within the study area. Impacts to barn swallow habitat may include the loss of foraging habitat, increased fragmentation and increased human interaction. Impacts from increased human presence are anticipated to be negligible given the

existing development surrounding the subject property and availability of suitable habitat within the greater study area.

Mitigation measures intended to prevent negative impacts to foraging barn swallow are presented in Section 7.

#### **6.1.1.2 Eastern Wood-Pewee**

Eastern wood-pewee (*Contopus virens*) is a small, avian insectivore listed as a species of special concern in Ontario. (COSEWIC, 2012a).

Historical occurrences have been reported for eastern wood-pewee within 1 km of the subject property; however, they were not observed during the breeding bird surveys or any other field investigations.

As eastern wood-pewee were not identified on-site during targeted breeding bird surveys or any other field investigation, the project is not anticipated to impact eastern wood-pewee or their habitat.

#### **6.1.1.3 Wood Thrush**

The wood thrush (*Hylocichla mustelina*) is a medium-sized songbird, listed as a species of special concern in Ontario.

Historical occurrences have been reported for wood thrush within 1 km of the subject property; however, they were not observed during breeding bird surveys or any other the field investigations.

As wood thrush were not identified on-site during targeted breeding bird surveys or any other field survey, the project is not anticipated to impact wood thrush of their habitat.

### **6.2 Fish Habitat**

According to the Provincial Planning Statement (MMAH, 2024), “development and site alteration shall not be permitted in fish habitat except in accordance with provincial and federal requirements.” Fish habitat as defined in the Fisheries Act (Canada, 1985) means “spawning grounds and nursery, rearing, food supply and migration areas on which fish depend directly or indirectly in order to carry out their life processes.”

As no in-water work is anticipated as part of the development, impacts are anticipated to be temporary and indirect in nature.

Potential indirect impacts to water quality and fish habitat from development can include increased overland flow and concomitant sediment transport caused by an increase in impervious surface

area, increased nutrient loading through both overland and subsurface pathways resulting from landscaping practices and septic leachate.

Mitigation measures, intended to protect fish habitat on-site are presented in Section 7.

### 6.2.1 Headwater Drainage Features

H1 had water conveyance throughout the first and second visits but was dry during the third visit. Due to H1 running along the agricultural field and originating as an agricultural ditch, it was determined to have limited riparian vegetation. In accordance with the TRCA/CVC guidance, the contributing (ephemeral) hydrology coupled with the limited riparian vegetation results in the determination that the above noted HDF requires **mitigation**.

H2 was channelized and had water conveyance throughout the first visit but was dry during the second and third visit. As such, it was determined to have limited hydrology. In conjunction with limited terrestrial habitat, **no management** is required.

H3 had water conveyance throughout the first and second visits but was dry during the third and was determined to have valuable hydrology. The feature provides contributing fish habitat, limited terrestrial habitat and important riparian habitat (scrubland). As such, **conservation** is required for H3.

H3A had water conveyance throughout the first and second visits but was dry during the third investigation. In conjunction with having no defined feature, H3 was determined to have valuable hydrology. As such they were determined to have contributing hydrology. The feature provides contributing fish habitat, limited terrestrial habitat, and important meadow riparian habitat. As such, **mitigation** is required for H3A.

H3B and H4 were channelized and had standing water during the first and second visit, but dry conditions during third site investigation. Both features provide contributing fish habitat, limited terrestrial habitat, and important scrubland riparian habitat. As such **conservation** is required for the feature.

As per the proposed development, H2, H3, H3A, and H3B will be removed. These features are identified as no management required, conservation and mitigation. These HDF segments designated conservation, and mitigation will be unable to be afforded complete mitigation measures as per the TCRA/CVC guidance. However, considering the that neither of these four features currently connects to downstream surface water features and provides limited functions, it is GEMTECs opinion that conservation and mitigation are not required for these segments.

Mitigation measures to protect baseflow conditions of the site and to replicate functions of the HDFs are provided below in Section 7.3.

### 6.3 Species at Risk

As outlined in the Endangered Species Act (Ontario, 2007), only species listed as threatened or endangered and their general habitat receive automatic protection. When a species-specific recovery strategy is developed, a specific habitat regulation will be established, which eventually replaces the automatic habitat protection. Species of special concern and their habitat do not receive protection under the ESA.

Potential impacts associated with the proposed project to threatened or endangered species identified as having a moderate or high potential to occur on-site in Section 4.7, are discussed on a species-by-species basis in the subsections below.

#### 6.3.1 Blanding's Turtle

Blanding's turtles (*Emydoidea blandingii*) have a highly domed, smooth black carapace with small, irregular tan or yellow flecking. The most distinctive characteristic of this species is the bright yellow chin and throat. Their hinged plastron is yellow with a large dark blotch in the corner of each scute, but may also be entirely black (Oldham and Weller, 2000).

In Canada, Blanding's turtles are found throughout southern and south-central Ontario from south of Manitoulin Island to western Quebec. In Ontario, Blanding's turtles are often observed utilizing eutrophic habitats with clear water (COSEWIC, 2005). This turtle species occurs primarily in shallow water; adults are generally found in open or partially vegetated sites, where as juveniles prefer areas that contain thick aquatic vegetation. Blanding's turtles are known to make large overland journeys between connected lakes, rivers, streams, marshes or ponds, upwards of 6 km in a single active season. Overwintering occurs in permanent pools that average about one metre in depth, or slow flowing streams (COSEWIC, 2005).

During the site investigation, Blanding's turtles were not detected on-site however the site is located within a greater area of known Blanding's turtle occurrences. Based on NHIC observation data the species has been observed within 1 km to the south, southwest, west, east, and southeast of the grid encompassing the site.

As outlined in the MNRF general habitat description for Blanding's turtle, Category 1 habitat is defined as "the nest and the area within 30 m of the nest or overwintering sites and the area within 30 m of the site", Category 2 habitat is defined as "the wetland complex (i.e. all suitable wetlands or waterbodies within 500 m of each other) that extends up to 2 km from an occurrence and the area within 30 m around those suitable wetlands or waterbodies" and Category 3 habitat is defined as "the area between 30 m and 250 m around suitable wetlands and waterbodies identified as Category 2, within 2 km of an occurrence." The MNRF general habitat description for Blanding's turtle is provided in Appendix E.

Blanding's turtle nests (Category 1 habitat) are created in open habitats with low vegetation cover, loose soils, and high sun exposure such as in forest clearings, meadows, shorelines, beaches

and gravel roads (Ontario, 2021) and (COSEWIC, 2016). Suitable Blanding's turtle overwintering habitat typically includes permanent bogs, fens, marshes, ponds, channels or other habitats with free (unfrozen) shallow water. Blanding's turtle may also hibernate within graminoid shallow marsh areas of larger marsh complexes by burying into substrates in areas of pooled water. Blanding's turtle may also overwinter in seasonal pools or small excavated areas with standing water (Ontario, 2021).

Suitable Category 2 habitat for Blanding's turtles during the active season includes a variety of permanent aquatic features including wetlands such as marsh, swamps, ponds, fens, bogs, slow-flowing streams, shallow bays of lakes or rivers, as well as graminoid shallow marsh and slough forest habitats that are adjacent to larger marsh complexes (Ontario, 2021). Suitable wetlands used during the active season are typically eutrophic (mineral or organic nutrient-rich), shallow with a soft substrate composed of decomposing materials, and often have emergent vegetation, such as water lilies and cattails (Ontario, 2021) and (COSEWIC, 2016).

Although wetlands and ponds are used as movement corridors when available, females can make extensive movements through upland habitat to access nesting sites (Ontario, 2021). Blanding's turtles also make regular overland movements between wetlands throughout the active season in order to access Category 1 and 2 habitats within their home range (Ontario, 2021). Category 3 habitat provides essential movement corridors of up to 500 m between wetlands, which will encompass the areas that are most likely to be used for overland movement (Ontario, 2021).

During the site investigation Blanding's turtles were not detected on-site. Review of NHIC occurrence data indicates the species has been observed within 1 km of the site.

As regulated Blanding's turtle habitat extends up to 2 km from an observation, based conservatively on the NHIC observation data, the site has the potential for Blanding's turtle or their habitat to occur on-site. However, based on field observations, none of the vegetation communities identified on-site provide suitable wetland habitat for regulated Blanding's turtle habitat. Furthermore, the lack of standing water within the on-site HDFs do not provide suitable permanent aquatic habitat (i.e. no surface water) to support Category 1 or Category 2 habitat. The watercourse along the southwest boundary may provide suitable Category 2 habitat, but does not provide suitable overwintering conditions for Category 1. Further the habitat within the watercourse is extremely limited in function, relatively low water levels, minimal in-stream vegetation and the watercourse runs between two agricultural fields. This watercourse is considered low suitability for regulated habitat. While Category 3 habitat would extend from the permanent watercourse, this habitat is also extremely limited in function and form for Blanding's turtle as there are no other suitable aquatic habitats on-site that a turtle would be migrating to (i.e. no ponds, wetlands, or other permanent aquatic features).

Based conservatively on the NHIC observation, a total of 0.24 ha of Category 2 and 4.96 ha of Category 3 habitat is present on-site. Category 2 and 3 Blanding's turtle habitat is illustrated on Figure A.4 in Appendix A.

The general habitat description for Blanding's turtle outlines that generally compatible activities include small-scale alterations to land cover that do not impede overland movements or impair nesting sites. No suitable areas of nesting are present on-site.

The proposed project has the potential to impact a small amount of low quality Category 2 and 3 habitat on-site. However, outside of construction the proposed development is not anticipated to negatively impact the use or function of migratory habitat, as Blanding's turtles will still be able to utilize the habitat for migration.

Potential impacts to individual Blanding's turtle are anticipated to be minimal and are limited to increased interactions with transient Blanding's turtles, particularly during migratory periods. Migration and dispersal take place after the start of the active season, following ice-off, and in September when turtles return to their overwintering habitat. Nesting typically takes place between late May to early July. Given the proposed development and minimal impact potential to Blanding's turtle and their habitat, it is GEMTEC's opinion that standard avoidance and mitigation measures will be sufficient to mitigate impacts of the proposed project and no ministry consultation is required.

General mitigation measures and best practices intended to Blanding's turtles from negative impacts are discussed in Section 7.

### **6.3.2 Eastern Red Bat**

Eastern red bat (*Lasiurus borealis*) is a medium-large sized (typically 10-17 g), insectivorous bat found in Ontario. The fur of an eastern red bat is usually orange, but can vary from yellowish-red to yellowish-grey, with white or white-tipped hairs (COSEWIC, 2023).

The eastern red bat is found throughout Canada (except Prince Edward Island), the United States, and northeast Mexico; with distribution uncommon west of the Western Cordillera. In Ontario, the species occurs throughout Ontario, appearing as far north as James Bay (COSEWIC, 2023).

Eastern red bats overwinter in warmer climates in the southern extent of the United States, typically beneath leaf litter (COSEWIC, 2023). In comparison to many other Ontario bat species, they do not overwinter in caves. During the spring and summer months, they typically utilize the foliage of trees and occasionally shrubs for roosting habitat, with a preference for roosting near the edge of the crown and at sufficient heights to prevent access from mammalian predators (COSEWIC, 2023).

Although the forest habitat on-site does not meet the requirements to support bat maternity colonies, given the availability of habitat on-site and within the study area, there is a potential for eastern red bat to occur on the property, primarily for foraging or non-maternal roosting. Impacts to eastern red bat are primarily associated with habitat loss, encroachment and increased wildlife-human interaction. Mitigation measures intended to protect eastern red bat from impacts of the proposed development are discussed in Section 6.

### 6.3.3 Eastern Small-footed Myotis

Eastern small-footed myotis (*Myotis leibii*) is the smallest (typically 3-5 g), insectivorous bat found in Ontario. The fur of an eastern small-footed myotis is golden-brown in colour, with a distinct black mask across the face. The eastern small-footed myotis is very similar in appearance to the little brown myotis and is distinguishable by their small foot and keeled calcar (Fraser, MacKenzie & Davy, 2007).

The eastern small-footed myotis is found throughout eastern North America. In Ontario, the species has been observed in the areas south of Lake Superior across to the Ontario-Quebec border (Humphrey, 2017).

Eastern small-footed myotis overwinter primarily in caves and abandoned mines with low humidity and temperatures and stable microclimates (Humphrey, 2017). In comparison to other Ontario bat species, they are able to tolerate much colder temperatures, drier conditions and draftier locations for hibernating (Humphrey, 2017). During the spring and summer months, they utilize a variety of habitats for roosting, including under rocks or rock outcrops, in buildings, under bridges, or in caves, mines or hollow trees (Ontario, 2021a).

Although the vegetation community on-site does not meet the requirements to support bat maternity colonies, given the availability of habitat and buildings within the study area, there is a potential for eastern small-footed myotis to occur on the property, primarily for foraging or non-maternal roosting. Impacts to eastern small-footed myotis are primarily associated with habitat loss, encroachment and increased wildlife-human interaction.

Mitigation measures intended to protect eastern small-footed myotis from impacts of the proposed development are discussed in Section 7.

### 6.3.4 Hoary Bat

Hoary bat (*Lasiurus cinereus*) is a large (typically 16-38 g), insectivorous bat found in Ontario and is the largest bat found in Canada. The fur of a hoary bat is dense and include a complex mixture of colors, ranging from light to dark brown, and have white tipped hairs on the dorsal and ventral sides (COSEWIC, 2023). The hoary bat is distinguishable by the large size and light yellow-brown fur on the head, throat, and anterior margins of the wings (COSEWIC, 2023).

The hoary bat range spans across all provinces and territories within Canada, all the states within the United States, and has a wide distribution throughout Mexico (COSEWIC, 2023). In Ontario, the hoary bat is found throughout the province, and has been observed north of James Bay (COSEWIC, 2023).

Hoary bats overwinter in warmer climates in the southern extent of the United States, typically beneath leaf litter (COSEWIC, 2023). In comparison to many other Ontario bat species, they do not overwinter in caves. During the spring and summer months, they typically utilize the foliage of trees and occasionally shrubs for roosting habitat, with a preference for roosting near the edge of the crown and at sufficient heights to prevent access from mammalian predators (COSEWIC, 2023).

Although the forest habitat on-site does not meet the requirements to support bat maternity colonies, given the availability of habitat on-site and within the study area, there is a potential for hoary bat to occur on the property, primarily for foraging or non-maternal roosting. Impacts to hoary bat are primarily associated with habitat loss, encroachment and increased wildlife-human interaction. Mitigation measures intended to protect hoary bat from impacts of the proposed development are discussed in Section 6.

#### **6.3.5 Little Brown Myotis**

Little brown myotis (*Myotis lucifugus*) is a small (typically 4-11 g), insectivorous bat. The fur of a little brown myotis is bi-coloured; fur is a glossy brown with a darker coloured base. The tragus of the little brown myotis is long and thin, with a rounded tip (Fraser, MacKenzie & Davy, 2007).

In Canada, little brown myotis' occur throughout all of the provinces and territories (except Nunavut), with its range extending south through the majority of the United States as well. In Ontario, the little brown myotis is widespread in southern Ontario and has been found as far north as Moose Factory and Favourable Lake (Ontario, 2021b).

Little brown myotis overwinter in caves and abandoned mines, they require highly humid conditions and temperatures that remain above the freezing mark (Ontario, 2021b). During the summer months, maternity colonies are often located in buildings or large-diameter trees. Little brown myotis roost in trees and buildings. Foraging occurs over water and along waterways, forest edges and in gaps in the forest. Open fields and clear-cuts are not typically utilized for foraging (COSEWIC, 2013).

Although the vegetation community on-site does not meet the requirements to support bat maternity colonies, given the availability of habitat and buildings within the study area, there is a potential for little brown myotis to occur on the property, primarily for foraging or non-maternal roosting. Impacts to little brown myotis are primarily associated with habitat loss, encroachment and increased wildlife-human interaction. Mitigation measures intended to protect little brown myotis from impacts of the proposed development are discussed in Section 7.

### 6.3.6 Silver-Haired Bat

Silver-haired bat (*Lasionycteris noctivagans*) is a medium-sized (typically 9-17 g), insectivorous bat. The fur is one of the darkest of all bats in Canada, with black skin membranes and black to dark brown fur (COSEWIC, 2023).

In North America, the silver-haired bat is widely distributed and spans from the southern extent of the Canadian provinces to east-central Mexico (COSEWIC, 2023). In Canada, the distribution spans from coast to coast, but appears to be uncommon in Atlantic Canada. In Ontario, the species occurs throughout Ontario, appearing as far north as James Bay (COSEWIC, 2023).

Silver-haired bat overwinter in mines, rock crevices, trees, and snags across North America, including the United States, the Great Lakes region of Ontario, and in some areas of British Columbia (COSEWIC, 2023). Foraging typically occurs in young and old forests. Silver-haired bat roost primarily under bark and in cavities of trees; however, may occasionally roost on or in buildings (COSEWIC, 2023).

Although the forest habitat on-site does not meet the requirements to support bat maternity colonies, given the availability of habitat and buildings on-site and within the study area, there is a potential for silver-haired bat to occur on the property, primarily for foraging or non-maternal roosting. Impacts to silver-haired bat are primarily associated with habitat loss, encroachment and increased wildlife-human interaction. Mitigation measures intended to protect silver-haired bat from impacts of the proposed development are discussed in Section 6.

### 6.3.7 Tri-colored Bat

Tri-colored bat (*Perimyotis subflavus*) is a small (typically 5-7 g), insectivorous bat. The fur is uniformly coloured on the ventral and dorsal sides, however, when parted fur shows three distinct colour bands. The base of the hair is blackish, with a blonde middle and brownish tip. The snout of the tri-coloured bat is also distinct, with swollen bulbous glands present (Fraser, MacKenzie & Davy, 2007).

In Canada, the tri-colored bat has only been recorded in southern parts of Nova Scotia, New Brunswick, Quebec and central Ontario. In Ontario, it occurs primarily from the southern edge of Lake Superior across to the Ontario-Quebec border and south (COSEWIC, 2013).

Tri-colored bat overwinter in caves or mines and have very rigid habitat requirements; they typically roost in the deepest parts where temperatures are the least variable and have the strongest correlation with humidity levels and warmer temperatures (COSEWIC, 2013). In the spring and summer, tri-colored bat utilizes trees, rock crevices and buildings for maternity colonies. Foraging is mainly done over watercourses and streamside vegetation (COSEWIC, 2013).

Although the vegetation community on-site does not meet the requirements to support bat maternity colonies, given the availability of habitat and buildings within the study area, there is a potential for tri-colored bat to occur on the property, primarily for foraging or non-maternal roosting. Impacts to tri-colored bat are primarily associated with habitat loss, encroachment and increased wildlife-human interaction. Mitigation measures intended to protect tri-colored bat from impacts of the proposed development are discussed in Section 7.

#### **6.4 Cumulative Impacts**

Potential cumulative impacts associated with the proposed project include an increase in storm water generation, increases in nutrient loading to adjacent aquatic features and the loss of forest habitat, primarily for avian species.

Cumulative impacts to the natural environment at the site due to increased human presence, wildlife and human interaction and increased noise, are expected to be negligible given the existing residential and major roadways surrounding the project area.

Cumulative impacts such as those listed above can be mitigated through development setbacks and recommended mitigation measures outlined in Section 7 below.

## 7.0 RECOMMENDED AVOIDANCE AND MITIGATION MEASURES

The following avoidance and mitigation measures have been recommended by GEMTEC in order to minimize or eliminate potential environmental impacts identified in Section 6.

For the purpose of this report, a setback is defined as the minimum required distance between any structure, development or disturbance and a specified line. A buffer, for the purpose of this report, is defined as the area located between a natural heritage feature and the prescribed setback. For the purpose of the following subsections, buffers should be located between natural heritage features and lands subject to development or alteration, be permanently vegetated by native or non-invasive, self sustaining vegetation and protect the natural heritage feature against the impact of the adjacent land use.

Vegetated buffers, particularly buffers that are vegetated with a mix of grassy herbaceous vegetation and shrubby or woody vegetation are most effective in mitigating impacts associated with anthropogenic activities in adjacent lands (Beacon, 2012). In the subsections below, where possible, literature references for studies used as the basis of the recommended buffer widths are provided.

### 7.1 Fish Habitat

Beacon Environmental Review of Ecological Buffers (2012), provides a range for buffer widths to protect various natural heritage features based on the current science. The buffers are presented in a way that determines the risk of not achieving the desired buffer function (i.e. high, moderate and low). The functions analysed include water quantity, water quality, screening or human disturbance/changes in land use, hazard mitigation zone and core habitat protection. Impacts to the watercourse on and off-site were identified to include potential impacts to water quality, human disturbance and core habitat protection (habitat for Blanding's turtle). Watercourse buffer widths have a moderate risk of not providing adequate mitigation for water quality and human disturbance/land use change impacts at widths between 11 m and 30 m. Watercourse buffer widths have a moderate risk of not providing adequate mitigation for core habitat protection at widths between 21 m and 60 m.

In consideration of the watercourse, and the nature of the proposed development, a minimum 30 m setback from the watercourse is recommended. The recommended 30 m setback provides sufficient protection for mitigating water quality impacts and human disturbances. At 30 m, the protection the buffer offers for core habitat protection, falls into the moderate risk of not achieving desired buffer function. As such a 30 m setback is sufficient to protect water quality impacts, human disturbance/land use change impacts, and core habitat within the watercourse. The setback is consistent with the recommendations of the Carp River Watershed/Subwatershed Study, which recommends a minimum 30 m setback. The 30 m setback is illustrated on Figure A.6 in Appendix A.

General mitigation measures recommended for the protection of water quality and fish habitat include:

- Buffers should be comprised of a mixture of native, self-sustaining trees, shrubs and tall grasses. The prescribed setbacks along the watercourse should remain in a natural, vegetated state.
- All future development and construction activities within the study area, including ditching, culvert installation, erosion and sediment control and storm water management should be completed in accordance with Ontario Provincial Standard Specification 182 and OPSS 805.
- Sediment fencing should be installed along all setbacks to provide visual demarcation of the setbacks to prevent machinery encroachment and sediment transport.
- Install and maintain effective sediment and erosion control measures before starting work.
- Schedule work to avoid wet, windy and rainy periods.
- When native soil is exposed, sediment and erosion control work in the form of heavy-duty sediment fencing shall be positioned along the down gradient edge of any construction envelopes adjacent to waterbodies.
- Maintain erosion and sediment control measures until all disturbed ground has been permanently stabilized, suspended sediment has resettled, and runoff water is clear.
- In order to protect fish habitat from contamination, it is recommended that all machinery be maintained in good working condition and that all machinery be fueled a minimum of 30 m from the high water mark.
- Any temporary storage of aggregate material shall be set back from the water's edge by no less than 40 m and be contained by heavy-duty silt fencing.
- Maintain as much permeable surface area as possible in future development plans to limit the generation of stormwater runoff.

### 7.1.1 Headwater Drainage Features

As detailed in Appendix D, a headwater drainage feature assessment was completed and six HDFs were identified on-site, identified as H1 through H4. Recommended management for on-site HDFs includes conservation, mitigation, and no management.

Following the classification guidance from TRCA/CVC, H3 and H3B is recommended for conservation management, and H3A is recommended for mitigation management. Conservation management typically requires maintaining, relocating and/or enhancing the existing features or riparian zone corridor; restoring lost functions through enhanced lot level controls; etc. and mitigation management typically requires replicating or enhancing feature functions through enhanced lot level conveyance measures (well vegetated swales), replicating on-site flow and outlet flow at the top of the system with vegetated swales/bioswales, restoring lost functions

through enhanced lot level controls and, replicating functions by lot level conveyance measures (e.g. vegetated swales) connected to the natural heritage system.

Currently, the existing features do not provide any direct connectivity to downstream surface water features, as such their function as an HDF from a landscape perspective is limited with respect to hydrology, terrestrial linkages, and fish habitat. As such it is GEMTECs opinion that the conservation management implications for H3 and H3B, and mitigation management implication for H3A do not accurately account for the limited connectivity of the features.

One of the key management implications for conservation and mitigation management features is to provide direct downstream connectivity. Currently neither features H2, H3, H3A or H3B are directly connected to downstream habitat. The proposed new drainage swales will collect surface run-off on site and feed it into the proposed SWM pond. The SWM pond will provide quality and quantity controls before discharging into the watercourse along the east property boundary. The new drainage swales and ditches will provide improved connectivity for surface water features on-site and off-site. The proposed SWM plan is sufficient to replicate and enhance the existing function of the HDFs on-site.

Where possible, drainage swales and ditches should be vegetated (i.e. bioswales) in order to enhance these features ability to attenuate nutrients and promote infiltration.

## **7.2 Significant Wildlife Habitat**

### **7.2.1 Habitats of Species of Conservation Concern - Barn Swallow**

Impacts to barn swallow from the proposed development primarily concern vegetation loss and increased fragmentation. As mentioned in Section 6.1.1.1, the site is likely to provide foraging habitat, however, nesting structures and an abundance of foraging habitat is present within the greater study area.

To minimize the impact of the proposed development on barn swallow, vegetation removal should occur outside the key breeding bird period (typically March 31 to August 31) as identified by Environment Canada for the protection of nesting and foraging barn swallow and to avoid contravention of the Migratory Bird Convention Act. If vegetation clearing activities must take place during the aforementioned timing window than a nest survey shall be conducted by a qualified professional.

## **7.3 Species at Risk**

### **7.3.1 Blanding's Turtle**

As discussed in Section 6.3.1, it is GEMTECs opinion that the proposed project will not negatively impact the function of regulated habitat on-site. As such it is GEMTECs opinion that standard avoidance and mitigation measures will be sufficient to mitigate impacts of the proposed project and no ministry consultation is required.

The 30 m setback from the permanent watercourse is sufficient to protect transient Blanding's turtles during migration. As mentioned in Section 6.3.1, approximately 0.24 ha of Category 2 Blanding's turtle habitat is present on-site, located within Block 7. Implementation of a 30 m setback from the watercourse, the potential loss of Category 2 habitat on-site is reduced from 0.24 ha to 0 ha on-site. Minor work within the Category 2 habitat is anticipated, to allow for the SWM pond outlet, however outside of construction of the pond and outlet, the proposed development is not anticipated to impact the availability or function of Category 2 habitat on-site. The 30 m setback ensures that riparian vegetation remains intact within the potential Category 2 habitat.

The following best practice measures should be implemented to avoid impacts to transient Blanding's turtles. If these measures cannot be implemented, consultation with the MECP is required to determine whether the project requires an authorization:

- Except for the SWM pond outlet, all future development, including the SWM pond, roadways and future buildings should be located outside of the 30 m setback. Minor grading to allow for the SWM pond outlet is permitted, but all other grading, site servicing and SWM components must occur outside of this setback.
- Prior to any site work, silt fencing should be installed around the entire construction area to prohibit the potential migration of Blanding's Turtles, and other wildlife into the construction area. Silt fencing should follow the protocols outlined in the Species at Risk Branch: Best Practices Technical Note: Reptile and Amphibian Exclusion Fencing Version 1.1 (MNRF, July 2013). Temporary fencing should be installed prior to the start of the active season and remain in place throughout the active season of each year of construction.
- Temporary exclusion fencing should be inspected by a designated staff member once per week between April 15 and October 15 of any year. The designated staff member should be trained by a Qualified Professional. Any damage to temporary fencing should be repaired by the end of the business day when the damage is observed.
- Each day of construction a daily pre-work sweep of the construction area should occur to ensure no SAR are present and to remove any wildlife from inside the construction area.
- All staff working on-site should be provided Species at Risk training to identify species at risk which a potential to occur on-site including: Blanding's turtle. Training will also outline the stop work procedures and MECP reporting/consultation prior to resuming work.
- During construction if any SAR is identified on-site all work should stop and a qualified professional and the MECP should be contacted for next steps. Sightings should be reported to the MECP and the NHIC.
- Heavy-duty silt fencing should be installed and maintained during construction and whenever soil is exposed.

- Tree clearing and vegetation removal will be undertaken outside of the active season (April 1 – October 31) for Blanding's turtles. Prior to vegetation removal a sweep will be completed to ensure Blanding's turtles are absent from the area.
- Cover all stock piled material with a geotextile to prevent turtles from nesting in the material between May 1 and August 1 of any year.
- To protect aquatic habitat for Blanding's turtles, machinery should be maintained in good working condition and all machinery should be fueled a minimum of 30 m from the high-water mark.
- To protect aquatic habitat, machinery should be maintained in good working condition and all machinery should be fueled a minimum of 30 m from the high-water mark.

### **7.3.2 Eastern Red Bat, Eastern Small-footed Myotis, Hoary Bat, Little Brown Myotis, Silver-Haired Bat & Tri-colored Bat**

As no critical habitat (i.e. overwintering caves or crevasses, or maternity roosts) were identified on-site, in accordance with MECP best management practices, to protect roosting and foraging bats, tree removal where required shall take place outside of the spring and summer active season (typically March 15 to November 30), when bats are more likely to be using forest habitat. If vegetation clearing cannot avoid the active season, the consultation with the MECP is needed to determine whether the project will require an authorization. If vegetation clearing must be conducted during the spring and summer timing window, then consultation with the MECP is required to determine whether the project will require an authorization.

To further protect bat species during vegetation removal, trees and vegetation (during the appropriate timing window) should be cleared in stages, working from the outer edge, in towards the centre, in order to provide wildlife in the forest time to migrate out.

In GEMTECs experience on similar development applications and consultation with the MECP for projects and properties of similar size and scale, the above mitigation/avoidance measures are sufficient to ensure no negative impacts to SAR bats. In eastern Ontario habitat is not a limiting factor, as such the MECP recommends the use of avoidance timing window for clearing of trees (>10cm in diameter) in order to avoid impacts to SAR bat species. As long as timing windows can be adhered to, the project will not impact SAR bats, and it is GEMTECs opinion that no further consultation with the MECP is required.

Should any components of the proposed project require tree clearing between March 15 and November 30, further consultation with the MECP is required.

## **7.4 Wildlife**

The following avoidance and mitigation measures are provided in effort to minimize impacts to on-site and off-site wildlife:

- Vegetation removal should occur outside of March 15 to November 30 to avoid the key breeding bird period, active turtle season, and bat summer active season. The timing window provides protection for migratory birds, SAR turtles, roosting bats and avoids contravention of the Migratory Bird Convention Act and Endangered Species Act. If vegetation clearing activities must take place during the aforementioned timing window then a nest survey and site sweep shall be conducted by a qualified professional to ensure no impacts to birds or turtles. If vegetation removal has the potential to impact SAR bats (i.e. vegetation removal within contiguous forested tracts) consultation with the MECP is required to determine whether the project will require an authorization.
- Installation of silt fence barriers around the entire construction envelope to prohibit the emigration of wildlife into the construction area, silt fencing should be checked daily and following each precipitation event.
- Cover all stockpiled material with a geotextile to prevent turtles from nesting in the material between May 1 and August 1 of any year.
- Perform daily pre-work sweeps of the construction area to ensure no species at risk are present and to remove any wildlife from inside the construction area.
- Should any species at risk be discovered throughout the course of the proposed works, the species at risk biologist with the local MECP district shall be contacted immediately and operations ceased to avoid any negative impacts to species at risk or their habitat until further direction is provided by the MECP.

## **7.5 Best Practice Measures for Mitigation of Cumulative Impacts**

The following best practice measures are provided for the mitigation of cumulative impacts resulting from general construction and development activities;

- To protect trees identified to be retained during construction, the Critical Root Zone (CRZ) should be identified and fenced. The CRZ is defined as 10 cm from the base of the tree for every centimetre in diameter of the tree trunk measured at breast height.
- Maintain as much permeable surface as possible in future development plans to minimize the generation of stormwater runoff.
- Silt fencing should be installed along all setbacks to provide visual demarcation of the setbacks and to prevent machinery encroachment and sediment transport.
- Erosion and sediment control measures should be maintained until all disturbed ground has been permanently stabilized.
- In effort to offset the effect of vegetation clearing, consideration should be given to landscape planting with native tree species indicative of the Great Lakes – St. Lawrence Forest Region, such as white cedar, white spruce, red maple, and red oak.

## 8.0 CONCLUSIONS

The proposed project supported by this EIS is the development of the approximately 23 ha property for future subdivision development. The development will result in the construction of twelve commercial/industrial lots, a road to provide access, and commercial/industrial buildings on each lot.

Based on the results of the impact analysis, impacts to the natural environment are anticipated to be minimal. Provided that mitigation measures recommended in Section 7 are implemented as proposed, no significant residual negative impacts are anticipated from the proposed future development.

Following review of the information pertaining to the natural heritage features of the site, the following general conclusions are provided by GEMTEC in regards to the Environmental Impact Statement.

- No significant negative impacts to natural heritage features identified on-site, including fish habitat, *candidate* significant wildlife habitat, habitat of species at risk, from future development are anticipated.
- The proposed project complies with the natural heritage policies of the Provincial Planning Statement.
- The proposed development complies with the natural heritage policies of the City of Ottawa Official Plan.

## 9.0 LIMITATION OF LIABILITY

This report and the work referred to within it have been undertaken by GEMTEC Consulting Engineers and Scientists Ltd (GEMTEC), and prepared for TLC Holdings Ltd. and is intended for the exclusive use of TLC Holdings Ltd.. This report may not be relied upon by any other person or entity without the express written consent of GEMTEC, TLC Holdings Ltd.. Nothing in this report is intended to provide a legal opinion.

The investigation undertaken by GEMTEC with respect to this report and any conclusions or recommendations made in this report reflect the best judgements of GEMTEC based on the site conditions observed during the investigations undertaken at the date(s) identified in the report and on the information available at the time the report was prepared.

This report has been prepared for the application noted and it is based, in part, on visual observations made at the site, all as described in the report. Unless otherwise stated, the findings contained in this report cannot be extrapolated or extended to previous or future site conditions, or portions of the site that were unavailable for direct investigation.

Should new information become available during future work or other studies, GEMTEC should be requested to review the information and, if necessary, re-assess the conclusions presented herein.

We trust this report provides sufficient information for your present purposes. If you have any questions concerning this report, please do not hesitate to contact our office.

Sincerely,



Emily Pentz, B.Sc.  
Junior Biologist



Taylor Warrington, B.Sc.  
Biologist

## 10.0 REFERENCES

- Beacon Environmental. 2012. Ecological Buffer Guideline Review – Prepared for Credit Valley
- Cadman M.D., D.A. Sutherland, G.G. Beck, D. Lepage, and A.R. Couturier. 2007. Atlas of the Breeding Birds of Ontario, 2001-2005. Bird Studies Canada, Environment Canada, Ontario Field Ornithologists, Ontario Ministry of Natural Resources, and Ontario Nature. Toronto.
- Canada, Government of (Canada). 1985. Fisheries Act. R.S.C. 1985, c. F-14.
- Chapman, L.J., and Putnam, D.F. 1984. The Physiography of Southern Ontario. Ontario Geological Survey, Special Volume 2.
- COSEWIC. 2023. COSEWIC assessment and status on the Hoary Bat *Lasiurus cinereus*, Eastern Red Bat *Lasiurus borealis* and Silver-Haired Bat, *Lasionycteris noctivagans*, in Canada. Committee on the Status of Endangered Wildlife in Canada. Ottawa. Xxi + 100pp.
- COSEWIC. 2013. COSEWIC assessment and status report on the Little Brown Myotis *Myotis lucifugus*, Northern Myotis *Myotis septentrionalis* and Tri-coloured Bat *Perimyotis subflavus* in Canada. Committee on the Status of Endangered Wildlife in Canada. Ottawa. xxiv + 93 pp.
- COSEWIC. 2012a. COSEWIC assessment and status report on the Eastern Wood-pewee *Contopus virens* in Canada. Committee on the Status of Endangered Wildlife in Canada. Ottawa. x + 39 pp.
- COSEWIC. 2012b. COSEWIC assessment and status report on the Wood Thrush *Hylocichla mustelina* in Canada. Committee on the Status of Endangered Wildlife in Canada. Ottawa. ix + 46 pp.
- Crins, J., William., P. A. Grey, P. W. Uhlig, and M.C. Wester. 2009. The Ecosystems of Ontario, Part I: Ecozones and Ecoregions. Ontario Ministry of Natural Resources, Peterborough, Ontario.
- Dobbyn, J.S. 1994. Atlas of the Mammals of Ontario. Federation of Ontario Naturalists, Toronto.
- Fraser E., MacKenzie, A., and Davy, C. 2007. Photo Field Guide to the Bats of Ontario. Published by St. Thomas Field Naturalists Club Incorporated.
- Humphrey, C. 2017. Recovery Strategy for the Eastern Small-footed Myotis (*Myotis leibii*) in Ontario. Ontario Recovery Strategy Series. Prepared for the Ontario Ministry of Natural Resources and Forestry, Peterborough, Ontario. Vii + 76 pp.
- Lee, H. T. 2008. Draft Southern Ontario Ecological Land Classification. Ministry of Natural Resources: London, Ontario.

Oldham, M.J and W.F. Weller. 2000. Ontario Herpetofaunal Atlas.

Ontario, Government of (Ontario). 2021a. Eastern small-footed Myotis. Viewed online: July 21, 2022. Available online: <https://www.ontario.ca/page/eastern-small-footed-myotis>

Ontario, Government of (Ontario). 2021b. Little Brown Myotis. Viewed online: July 21, 2022. Available online: <https://www.ontario.ca/page/little-brown-myotis>

Ontario, Government of (Ontario). 2021c. Northern Myotis. Viewed online: July 21, 2022. Available online: <https://www.ontario.ca/page/northern-myotis>

Ontario, Government of (Ontario). 1990. Conservation Authorities Act. R.S.O. 1990. Chapter C.27. Last amendment: 2011, C.9 Sched. 27, S. 22.

Ontario Legislative Assembly. 2007. Endangered Species Act.

Ontario Ministry of Municipal Affairs and Housing (MMAH). 2024. Provincial Planning Statement – Under Planning Act, Toronto. October.

Ontario Ministry of Natural Resources and Forestry (OMNRF). 2018. Natural Heritage Information Request Guide.

Ontario Ministry of Natural Resources and Forestry (OMNRF). 2015. Significant Wildlife Habitat Ecoregion 6E Criterion Schedules.

Ontario Ministry of Natural Resources and Forestry (OMNRF). 2014a. Make a Map: Natural Heritage Areas.

Ontario Ministry of Natural Resources and Forestry (OMNRF). 2014b. Significant Wildlife Habitat Mitigation Support Tool.

Ontario Ministry of Natural Resources and Forestry (OMNRF). 2013. Natural Heritage Information Centre (NHIC) Biodiversity Explorer.

Ontario Ministry of Natural Resources and Forestry (OMNRF). 2011. Land Information Ontario (LIO).

Ontario Ministry of Natural Resources (OMNR). March 2010. Natural Heritage Reference Manual for Natural Heritage Policies of the Provincial Policy Statement, 2005. Second Edition.

Ontario Ministry of Natural Resources (OMNR). 2005 Natural Heritage Information Centre. Ontario Odonata Atlas.

Ontario Ministry of Natural Resources (OMNR). 2000. Significant Wildlife Technical Guide.

Ontario Geological Survey 2019. Surficial geology of Southern Ontario; Ontario Geological Survey, Miscellaneous Release--Data 128-REV

Ontario Nature, 2019, Ontario Reptile and Amphibian Atlas. Viewed online: Viewed online November 24, 2023. Available online:

<https://www.ontarioinsects.org/herp/index.html?Sort=1&area2=squaresCounties&records=all&myZoom=5&Lat=42.95&Long=-81.01>

Ottawa, City of (Ottawa). 2023, Environmental Impact Study Guidelines.

Ottawa, City of (Ottawa). 2022a. Bird-Safe Design Guidelines. December 2022.

Ottawa, City of (Ottawa). 2022b. City of Ottawa Official Plan.

Ottawa, City of (Ottawa). 2022c. City of Ottawa Protocol for Wildlife Protection During Construction. August 2015.

Ottawa, City of (Ottawa). 2022d. Significant Woodlands: Guidelines for Identification, Evaluation, and Impact Assessment. Draft.

Robinson Consultants Inc. (2004). Carp River Watershed/ Subwatershed Study. Volume I – Main Report. December 2004.



## **APPENDIX A**

### Report Figures

Figure A.1 – Site Location

Figure A.2 – Site Layout

Figure A.3 – Vegetation Communities

Figure A.4 – Natural Heritage Features


Figure A.5 – Proposed Development

Figure A.6 - Mitigation Measures



**Legend**

- Property Boundary
- Study Area



 <p><b>GEMTEC</b> CONSULTING ENGINEERS AND SCIENTISTS</p>	32 Steacie Drive, Ottawa, ON K2K 2A9 T: (613) 836-1422 www.gemtec.ca ottawa@gemtec.ca
--	---


Client: <b>TLC Holdings Ltd.</b>	Project: <b>102151.001</b>
----------------------------------	----------------------------

Location	<b>3160 Carp Road Ottawa, Ontario</b>
----------	---

Drwn By: EP	Chkd By: TW	<b>Site Location</b>
----------------	----------------	----------------------


Date: December 2024	Rev. 0	<b>Figure: A.1</b>
© Queen's Printer for Ontario		

Coordinate System: NAD 1983 UTM Zone 18N  
 Service Layer Credits: World Street Map: City of Ottawa, Province of Ontario, Esri Canada, Esri, TomTom, Garmin, SafeGraph, GeoTechnologies, Inc, METI/NASA, USGS, EPA, NPS, US Census Bureau, USDA, NRCAN, Parks Canada  
 World Topographic Map: City of Ottawa, Province of Ontario, Esri Canada, Esri, TomTom, Garmin, SafeGraph, METI/NASA, USGS, EPA, NPS, USDA, NRCAN, Parks Canada



### Legend

- Property Boundary
- Study Area
- Watercourse
- Breeding Bird Survey Location (100 m Radius)
- - - - ● Headwater Drainage Feature



**GEMTEC**  
CONSULTING ENGINEERS  
AND SCIENTISTS

32 Steacie Drive,  
Ottawa, ON K2K 2A9  
T: (613) 836-1422  
www.gemtec.ca  
ottawa@gemtec.ca

Client: <b>TLC Holdings Ltd.</b>	Project: 102151.001
----------------------------------	---------------------

Location  
**3160 Carp Road  
Ottawa, Ontario**

Drwn By: EP	Chkd By: TW	<b>Survey Locations</b>
----------------	----------------	-------------------------

Date: December 2024	Rev. 0	<b>Figure: A.2</b>
© Queen's Printer for Ontario		


Coordinate System: NAD 1983 UTM Zone 18N  
 Service Layer Credits: Hybrid Reference Layer: Esri Community Maps Contributors, City of Ottawa, Province of Ontario, Esri Canada, Esri, TomTom, Garmin, SafeGraph, GeoTechnologies, Inc, METI/NASA, USGS, EPA, NPS, US Census Bureau, USDA, NRCan, Parks Canada  
 World Imagery: Maxar



**Legend**

- Property Boundary
- Study Area
- Watercourse
- - --- - ● Headwater Drainage Feature
- Vegetation Community

FODM7 = Fresh – Moist Lowland Deciduous Forest  
 THDM2-6 = Buckthorn Deciduous Shrub Thicket  
 ME = Meadow  
 MEM = Mixed Meadow  
 OAG = Open Agriculture



**GEMTEC**  
CONSULTING ENGINEERS  
 AND SCIENTISTS

32 Steacie Drive,  
 Ottawa, ON K2K 2A9  
 T: (613) 836-1422  
 www.gemtec.ca  
 ottawa@gemtec.ca

Client: <b>TLC Holdings Ltd.</b>	Project: 102151.001
-------------------------------------	------------------------

Location  
**3160 Carp Road  
 Ottawa, Ontario**

Drwn By: EP	Chkd By: TW	<b>Vegetation Communities</b>
----------------	----------------	-------------------------------


Date: December 2024	Rev. 0	<b>Figure: A.3</b>
© Queen's Printer for Ontario		

Coordinate System: NAD 1983 UTM Zone 18N  
 Service Layer Credits: Hybrid Reference Layer: Esri Community Maps Contributors, City of Ottawa, Province of Ontario, Esri Canada, Esri, TomTom, Garmin, SafeGraph, GeoTechnologies, Inc, METI/NASA, USGS, EPA, NPS, US Census Bureau, USDA, NRCan, Parks Canada  
 World Imagery: Maxar



**Legend**

- Property Boundary
- Study Area
- Watercourse
- Headwater Drainage Feature
- Candidate Blanding's Turtle Habitat 2 (30 m)
- Candidate Blanding's Turtle Habitat 3 (250 m)



**GEMTEC**  
CONSULTING ENGINEERS  
AND SCIENTISTS

32 Steacie Drive,  
Ottawa, ON K2K 2A9  
T: (613) 836-1422  
www.gemtec.ca  
ottawa@gemtec.ca

Client: <b>TLC Holdings Ltd.</b>	Project: 102151.001
-------------------------------------	------------------------

Location  
**3160 Carp Road  
Ottawa, Ontario**

Drwn By: EP	Chkd By: TW	<b>Natural Heritage Features</b>
----------------	----------------	----------------------------------

Date: December 2024	Rev. 0	<b>Figure: A.4</b>
© Queen's Printer for Ontario		

Coordinate System: NAD 1983 UTM Zone 18N  
 Service Layer Credits: Hybrid Reference Layer: Esri Community Maps Contributors, City of Ottawa, Province of Ontario, Esri Canada, Esri, TomTom, Garmin, SafeGraph, GeoTechnologies, Inc, METI/NASA, USGS, EPA, NPS, US Census Bureau, USDA, NRCan, Parks Canada  
 World Imagery: Maxar




**Legend**

- Property Boundary
- Study Area
- Watercourse
- Headwater Drainage Feature
- Candidate Blanding's Turtle Habitat 2 (30 m)
- Candidate Blanding's Turtle Habitat 3 (250 m)

**Development Concept**

- Proposed Lot
- Proposed Road



**GEMTEC**  
CONSULTING ENGINEERS  
AND SCIENTISTS

32 Steacie Drive,  
Ottawa, ON K2K 2A9  
T: (613) 836-1422  
www.gemtec.ca  
ottawa@gemtec.ca


Client: <b>TLC Holdings Ltd.</b>	Project: 102151.001
----------------------------------	---------------------

Location  
**3160 Carp Road  
Ottawa, Ontario**

Drwn By: EP	Chkd By: TW	<b>Proposed Development</b>
----------------	----------------	-----------------------------

Date: December 2024	Rev. 0	<b>Figure: A.5</b>
© Queen's Printer for Ontario		

Coordinate System: NAD 1983 UTM Zone 18N  
 Service Layer Credits: Hybrid Reference Layer: Esri Community Maps Contributors, City of Ottawa, Province of Ontario, Esri Canada, Esri, TomTom, Garmin, SafeGraph, GeoTechnologies, Inc, METI/NASA, USGS, EPA, NPS, US Census Bureau, USDA, NRCAN, Parks Canada  
 World Imagery: Maxar




**Legend**

- Property Boundary
- Study Area
- Watercourse
- Candidate Blanding's Turtle Habitat 3 (250 m)
- Candidate Blanding's Turtle Habitat 2 (30 m)
- Development Concept
- 30 m Setback

**Headwater Drainage Feature Management**

- Conservation
- Mitigation
- No Management
- Proposed Realigned Headwater Drainage Features



**GEMTEC**  
CONSULTING ENGINEERS  
AND SCIENTISTS

32 Steacie Drive,  
Ottawa, ON K2K 2A9  
T: (613) 836-1422  
www.gemtec.ca  
ottawa@gemtec.ca

Client: <b>TLC Holdings Ltd.</b>	Project: 102151.001
----------------------------------	---------------------

Location  
**3160 Carp Road  
Ottawa, Ontario**

Drwn By: EP	Chkd By: TW	<b>Mitigation Measures</b>
----------------	----------------	----------------------------

Date: March 2025	Rev. 1	<b>Figure: A.6</b>
© Queen's Printer for Ontario		

Coordinate System: NAD 1983 UTM Zone 18N  
Service Layer Credits: World Imagery: Maxar  
Hybrid Reference Layer: Sources: Esri, TomTom, Garmin, FAO, NOAA, USGS, © OpenStreetMap contributors, and the GIS User Community




## **APPENDIX B**

Site Photographs



Site Photograph 1: Mixed Meadow (MEM).



Site Photograph 2: Fresh – Moist Lowland Deciduous Forest (FODM7).



Site Photograph 3: Open Agriculture (OAG).



Site Photograph 4: Meadow (ME).



Site Photograph 5: Upstream extent of HDF3 and Buckthorn Deciduous Shrub Thicket (THDM2-6).




Site Photograph 6: The vertical drain on-site.




Site Photograph 7: HDF1



Site Photograph 8: HDF2



Site Photograph 9: Downstream extent of HDF3.




Site Photograph 10: HDF3A



Site Photograph 11: HDF3B



Site Photograph 12: HDF4



Site Photograph 12: Watercourse along east property boundary.



## **APPENDIX C**

Report Summary Tables

**TABLE C.1  
SUMMARY OF WILDLIFE OBSERVED ON-SITE AND ADJACENT TO SITE**

Common Name	Scientific Name	S-Rank	Evidence
<b>Avian Species</b>			
American crow	<i>Corvus brachyrhynchos</i>	S5	Heard calling
American goldfinch	<i>Spinus tristis</i>	S5	Heard calling
American robin	<i>Turdus migratorius</i>	S5	Heard calling
Barn swallow	<i>Hirundo rustica</i>	S4B	Observed flying overhead, foraging
Black-capped chickadee	<i>Poecile atricapillus</i>	S5	Heard calling
Blue jay	<i>Cyanocitta cristata</i>	S5	Heard calling
Brown thrasher	<i>Toxostoma rufum</i>	S4B	Heard calling
Canada goose	<i>Branta canadensis</i>	S5	Observed
Cedar waxwing	<i>Bombycilla cedrorum</i>	S5	Heard calling
Chestnut-sided warbler	<i>Setophaga pensylvanica</i>	S5B	Heard calling
Common grackle	<i>Quiscalus quiscula</i>	S5	Heard calling
Common yellowthroat	<i>Geothlypis trichas</i>	S5B,S3N	Heard calling
Gray catbird	<i>Dumetella carolinensis</i>	S5B,S3N	Heard calling
Great crested flycatcher	<i>Myiarchus crinitus</i>	S5B	Heard calling
Killdeer	<i>Charadrius vociferus</i>	S4B	Heard calling
Mourning dove	<i>Zenaida macroura</i>	S5	Heard calling
Northern cardinal	<i>Cardinalis cardinalis</i>	S5	Heard calling
Northern flicker	<i>Colaptes auratus</i>	S5	Heard calling
Red-eyed vireo	<i>Vireo olivaceus</i>	S5B	Heard calling
Red-winged blackbird	<i>Agelaius phoeniceus</i>	S5	Heard calling
Ring-billed gull	<i>Larus delawarensis</i>	S5	Heard calling
Rose-breasted grosbeak	<i>Pheucticus ludovicianus</i>	S5B	Heard calling
Song sparrow	<i>Melospiza melodia</i>	S5	Heard calling
Turkey vulture	<i>Cathartes aura</i>	S5B,S3N	Observed flying overhead
Veery	<i>Catharus fuscescens</i>	S5B	Heard calling
White-throated sparrow	<i>Zonotrichia albicollis</i>	S5	Heard calling
Wild turkey	<i>Meleagris gallopavo</i>	S5	Observed
Yellow warbler	<i>Setophaga petechia</i>	S5B	Heard calling
<b>Mammalian Species</b>			
White-tailed deer	<i>Odocoileus virginianus</i>	S5	Observed on-site

**Notes:**

\* Denotes a Species at Risk

Subnational Conservation Status Ranks:

S1 - Critically Imperilled, at very high risk of extirpation, very few populations or occurrences or very steep population decline

S2 - Imperiled, at high risk of extirpation, few populations or occurrences or steep population decline

S3 - Vulnerable, at moderate risk of extirpation, relatively few populations or occurrences, recent and widespread population decline

S4 - Apparently Secure, at a family low risk of extirpation, many populations or occurrences, some concern for local population decline

S5 - Secure, at very low or no risk of extirpation, abundant populations or occurrences, little to no concern for population decline

Qualifiers:

S#B - Conservation status refers to the breeding population of the species

S#N - Conservation status refers to the non-breeding population of the species

S#M - Migrant species, conservation status refers to the aggregating transient population of the species

**TABLE C.2  
SCREENING RATIONAL FOR SIGNIFICANT WOODLANDS**

Woodland Criteria	Further Considered in EIS	Rationale
Woodland Size	No	Contiguous woodlands on-site do not meet the minimum size requirement for the planning area (>50 ha).
Ecological Functions		
a) Woodland Interior	No	No interior woodlands present on-site.
b) Proximity	No	The woodlands on-site are not within proximity to other natural heritage features.
c) Linkages	No	The woodlands on-site do not provide linkages to other natural heritage features.
d) Water Protection	Yes	Woodlands on-site are proximate to fish habitat.
e) Diversity	No	Species composition within the on-site woodland is well represented on the landscape and no rare species communities were observed on-site.
Uncommon Characteristics	No	The woodlands on-site do not have a unique species composition, vegetation communities with a ranking of S1, S2 or S3, or a mature size structure.
Economical and Social Functional Values	No	The woodlands on-site do not contain high productivity in terms of economically valuable products, high social value such as recreational use, identified historical cultural or educational values.

**TABLE C.3  
SCREENING RATIONALE FOR HABITATS OF SEASONAL CONCENTRATION AREAS**

<b>Wildlife Habitat</b>	<b>Further Considered in EIS</b>	<b>Rationale</b>
Waterfowl Stopover and Staging Areas	No	No suitable habitat located on-site or within the study area to support waterfowl stopover and staging areas.
Shorebird Migratory Stopover Area	No	Shorebird stopover sites are typically well-known and have a long history of use. The site does not contain suitable shoreline habitat for shorebird foraging.
Raptor Wintering Area	No	No suitable woodland and upland habitat to support raptor overwintering.
Bat Hibernacula	No	Cave and crevice habitat is not present on-site or within the study area.
Bat Maternity Colonies	No	Woodlands on-site do not meet minimum snag density (>10 snags/hectare) requirement to be considered SWH for bat maternity colonies.
Turtle Wintering Area	No	No suitable habitat on-site or within the study area to support turtle overwintering.
Reptile Hibernaculum	No	No structures such as large rock piles, bedrock outcrops, cervices or other karstic features have been identified on-site.
Colonial Bird Nesting Habitat	No	No suitable habitat located on-site or within the study area to support colonial bird nesting.
Migratory Butterfly Stopover Area	No	The site is not located within 5 km of Lake Ontario and therefore does not meet the defining criteria.
Landbird Migratory Stopover Area	No	The site is not located within 5 km of Lake Ontario and therefore does not meet the defining criteria.
Deer Yarding Areas and Winter Congregation Areas	No	As outlined in the Significant Wildlife Habitat Criteria Schedules (OMNRF, 2015) winter deer yards and deer management are an MNRF responsibility. Based on review of publicly available data from the OMNRF on Land Information Ontario Geo-hub, no Stratum I deer yards, Stratum II deer yards, or winter congregation areas have been identified on-site or within the broader study area.

**TABLE C.4**  
**SCREENING RATIONALE FOR SPECIALIZED WILDLIFE HABITATS**

Specialized Wildlife Habitat	Further Considered in EIS	Rationale
Waterfowl Nesting Area	No	No upland habitat adjacent to wetland is present on-site to support waterfowl nesting areas.
Bald Eagle and Osprey Nesting, Foraging and Perching Habitat	No	The site is located >120 m from any habitat which could support foraging bald eagles or osprey. Nesting sites for these species are uncommon in Ecoregion 6E (MNRF, 2012).
Woodland Nesting Raptor Habitat	No	Nesting may occur in any ecosite and species preference is towards mature forest stands >30 ha with >10 ha of interior habitat with a 200 m buffer. Contiguous forest stands >30 ha are not present on-site. No sticks nests were observed on-site.
Turtle Nesting Habitat	No	No wetland on-site to provide suitable nesting habitat.
Seeps and Springs	No	No seeps or springs are present on-site.
Woodland Amphibian Breeding Habitat	No	No suitable wetland habitat adjacent to woodlands to support woodland amphibian breeding habitat.
Wetland Amphibian Breeding Habitat	No	No wetlands on-site or within study area to support wetland amphibian breeding habitat.
Woodland Area-Sensitive Bird Breeding Habitat	No	Woodland area-sensitive birds require interior forest habitat located >200 m from the forest edge in large (>30 ha) forest stands. Woodlands on-site and adjacent to the site do not meet the defining criteria.

**TABLE C.5  
SCREENING RATIONALE FOR HABITAT FOR SPECIES OF CONSERVATION CONCERN**

General Habitats of Species of Conservation Concern	Further Considered in EIS	Rationale
Marsh Breeding Bird Habitat	No	Suitable marsh habitat is not present on-site.
Open Country Breeding Bird Habitat	No	Candidate open country breeding bird habitat typically includes large grassland areas (includes natural and cultural fields and meadows) that are >30 ha. The meadow habitat on-site does not meet the size requirement; thus, no suitable habitat is present on-site.
Shrub/Early Successional Breeding Bird Habitat	No	Candidate early successional breeding bird habitat typically includes fallow fields transitioning to early successional forest habitats that are > 10 ha but have not been actively used for farming. No suitable habitat is present on-site.
Terrestrial Crayfish Habitat	No	Terrestrial crayfish are only found within southwestern Ontario (MNRF, 2012).
Special Concern and Rare Wildlife Species	Yes	One species of special concern were identified on-site during the site investigation: barn swallow. Occurrence data from the NHIC indicates historical occurrence of eastern wood-pewee and wood thrush within 1km of the site.

**TABLE C.6  
SCREENING RATIONALE FOR ANIMAL MOVEMENT CORRIDORS**

General Habitats of Species of Conservation Concern	Further Considered in EIS	Rationale
Amphibian Movement Corridor	No	No amphibian movement corridors have been identified on-site during the site investigation, nor has it been identified by MNR mapping. Additionally, no wetlands have been identified on site or within the study area.
Deer Movement Corridor	No	No cervid movement corridors have been identified on-site during the site investigation, nor has it been identified by MNR mapping.

**TABLE C.7  
SCREENING RATIONALE FOR POTENTIAL SPECIES AT RISK ON-SITE OR WITHIN STUDY AREA**

Species	ESA Status	Habitat Use	Probability of Occurrence On-Site or Within Study Area	Rationale
<b>Avian</b>				
Bank Swallow	Threatened	Colonial nester, burrows in eroding silt, to sand banks, sand pit walls, etc.	Low	NHIC data indicates bank swallow within 1km of the property. Site lacks suitable nesting habitat; however, it may be present in the greater study area. Species not observed on-site.
Barn Swallow	Special Concern	Nests in barns and other semi-open structures. Forages over open fields and meadows.	High	Site lacks suitable nesting habitat; however it may be present in the study area. Suitable foraging habitat is likely present on-site. Species observed during site investigations.
Bobolink	Threatened	Nests in dense tall grass fields and meadows, low tolerance for woody vegetation.	Low	Suitable habitat not present on-site. No historical occurrences recorded. Species not observed on-site.
Canada Warbler	Special Concern	Prefers wet forests with dense shrub layers	Low	Forest composition may provide suitable habitat. No historical records of species in study area. Species was not observed or detected during any of the site investigations.
Cerulean Warbler	Threatened	Prefers mature deciduous forest habitat.	Low	Forest structure on-site is unlikely to provide preferred habitat. Species was not observed or detected during any of the site investigation.
Chimney Swift	Threatened	Nests in traditional-style open brick chimneys.	Low	No anthropogenic structures on-site to provide preferred habitat; however, may occur within the study area.
Common Nighthawk	Special Concern	Nests in a variety of open sites: beaches, fields and gravel rooftops.	Low	Suitable habitat does not occur on-site. No historical occurrences recorded. Species not observed on-site.
Eastern Meadowlark	Threatened	Nests and forages in dense tall grass fields and meadows, higher tolerance to woody vegetation.	Low	Suitable habitat not present on-site. No historical occurrences recorded. Species not observed on-site.
Eastern Whip-poor-will	Threatened	Nests on the ground in open deciduous or mixed woodlands with little underbrush, and bedrock outcrops.	Low	No suitable habitat may be present on-site or within the study area. No historical occurrences recorded. Species not observed on-site.
Eastern Wood-Pewee	Special Concern	Woodland species, often found near clearings and edge habitat.	Moderate	NHIC data indicates eastern wood-pewee historical occurrences within 1km from the site. Suitable forest habitat may occur on-site. Species not observed during the site investigations.
Evening Grosbeak	Special Concern	Nests in trees or large shrubs, preference to large coniferous forests, will use deciduous. Overwinters in Ottawa.	Low	Forest structure on-site is unlikely to provide preferred habitat. Species was not observed or detected during any of the site investigation.
Golden Eagle	Endangered	Nests on remote, bedrock cliffs, overlooking large burns, lakes or tundra	Low	Suitable nesting habitat does not occur on-site. No historical occurrences recorded. Species not observed on-site.
Golden-winged Warbler	Special Concern	Ground nesting, edge species. Breeds in successional scrub habitats surrounded by forests.	Low	Site is unlikely to provide suitable habitat for golden-winged warblers due to the lack of successional scrub habitat.
Grasshopper Sparrow	Special Concern	Ground-nesting grassland species. Prefers fields with low sparse vegetation on sand, alvars or poor soils.	Low	Suitable habitat does not occur on-site. No historical occurrences recorded. Species not observed on-site.
Henslow's Sparrow	Endangered	Prefers open, moist, tallgrass fields.	Low	Suitable habitat does not occur on-site. No historical occurrences recorded. Species not observed on-site.
Least Bittern	Threatened	Prefers marshes, shrub swamps, usually near cattails	Low	Suitable wetland habitat does not occur on-site. No historical occurrences recorded. Species not observed on-site.
Loggerhead Shrike	Endangered	Prefers grazed pastures with short grass and scattered shrubs, especially hawthorn.	Low	Suitable habitat does not occur on-site. No historical occurrences recorded. Species not observed on-site.
Olive-sided Flycatcher	Special Concern	Forest edge species, forages in open areas from high vantage points in trees.	Low	Suitable forest habitat does not occur on-site. No historical occurrences recorded. Species not observed on-site.
Peregrine Falcon	Special Concern	Nests on cliffs near water and on more anthropogenic structures such as tall buildings, bridges, and smokestacks.	Low	Suitable nesting habitat does not occur on-site. No historical occurrences recorded. Species not observed on-site.
Red-headed Woodpecker	Endangered	Prefers open deciduous woodlands, particularly those dominated by oak and beech.	Low	Suitable forest habitat does not occur on-site. No historical occurrences recorded. Species not observed on-site.
Rusty Blackbird	Special Concern	Wet wooded or shrubby areas (nests at edges of Boreal wetlands)	Low	Suitable nesting habitat does not occur on-site. No historical occurrences recorded. Species not observed on-site.
Short-eared Owl	Threatened	Ground nester, prefers open habitats, fields and marshes.	Low	Suitable nesting habitat does not occur on-site. No historical occurrences recorded. Species not observed on-site.
Wood Thrush	Special Concern	Prefers deciduous or mixed woodlands.	Moderate	NHIC data indicates wood thrush historical occurrences within 1km from the site. Suitable forest habitat may occur on-site. Species not observed during the site investigations.

**TABLE C.7  
SCREENING RATIONALE FOR POTENTIAL SPECIES AT RISK ON-SITE OR WITHIN STUDY AREA**

Species	ESA Status	Habitat Use	Probability of Occurrence On-Site or Within Study Area	Rationale
<b>Mammalian</b>				
Eastern small-footed Myotis	Endangered	Roosts in rock crevices, barns and sheds. Overwinters in abandoned mines. Summer habitats are poorly understood in Ontario, elsewhere prefers to roost in open, sunny rocky habitat and occasionally in buildings (Humphrey, 2017).	Moderate	Available habitat within project area unlikely to meet bat maternity colony requirements however the site and surrounding area may provide foraging and non-maternal roost habitat.
Little Brown Myotis	Endangered	Maternal colonies known to use buildings, may also roost in trees during summer. Affinity towards anthropogenic structures for summer roosting habitat and exhibit high site fidelity (Environment Canada, 2015).	Moderate	Available habitat within project area unlikely to meet bat maternity colony requirements however the site and surrounding area may provide foraging and non-maternal roost habitat.
Northern myotis (Northern Long-eared Bat)	Endangered	Occurs throughout eastern North America in associated with Boreal forests. Roosts mainly in trees, occasionally anthropogenic structures during summer (Environment Canada, 2015). Overwinters in caves and abandoned mines.	Low	Species affinity is for Boreal forests and rarely roosts in anthropogenic structures. Subject property occurs at extreme southern end of species range and lacks preferred boreal forest habitat.
Tri-colored Bat	Endangered	Roosts in trees, rock crevices and occasionally buildings during summer. Overwinters in caves and mines.	Moderate	Available habitat within project area unlikely to meet bat maternity colony requirements however the site and surrounding area may provide foraging and non-maternal roost habitat.
<b>Reptilian</b>				
Blanding's Turtle	Threatened	Inhabits quiet lakes, streams and wetlands with abundant emergent vegetation. Frequently occurs in adjacent upland forests.	Moderate	NHIC observations indicated species has been observed within 1km of the site, however the site and study area do not provide suitable wetland habitat to support Blanding's turtle or their habitat. Occurrence records likely associated with wetland habitat south of site. Potential for transient Blanding's turtles to occur on-site.
Eastern Musk Turtle	Special Concern	Wetlands. Highly aquatic habitats.	Low	No historical occurrences for the species within the study area. No suitable aquatic habitat on-site. Species not observed on-site.
Eastern Ribbonsnake	Special Concern	Marshy edges of wetlands and watercourses.	Low	No historical occurrences for the species within the study area. No suitable aquatic habitat on-site. Species not observed on-site.
Northern Map Turtle	Special Concern	Highly aquatic species, found only in lakes and large rivers.	Low	No historical occurrences for the species within the study area. No suitable aquatic habitat on-site. Species not observed on-site.
Snapping Turtle	Special Concern	Highly aquatic species, found in a wide variety of wetlands, water bodies and watercourses.	Low	No historical occurrences for the species within the study area. No suitable aquatic habitat on-site. Species not observed on-site.
Spotted Turtle	Endangered	Secretive wetland species.	Low	No historical occurrences for the species within the study area. No suitable aquatic habitat on-site. Species not observed on-site.
Wood Turtle	Endangered	Primarily terrestrial forest species. Associated with clear, gravelly streams.	Low	No historical occurrences for the species within the study area. No suitable habitat on-site. Species not observed on-site.
<b>Plants</b>				
American Ginseng	Endangered	Rich, moist, relatively mature deciduous forests.	Low	Forest on-site is unlikely to provide suitable habitat.
Black Ash	Endangered	Predominantly a wetland species, found in swamps, floodplains and fens.	Low	No suitable habitat on-site to support species.
Butternut	Endangered	Inhabits a wide range of habitats including upland and lowland deciduous and mixed forests.	Low	Species not observed on-site.
<b>Lichens</b>				
Pale-bellied Frost Lichen	Endangered	Grows on the bark of hardwood trees such as white ash, black walnut, American elm and ironwood. Can also be found growing on fence posts and boulders.	Low	Species believed to be extirpated from the Ottawa area.
<b>Fish</b>				
American Eel	Endangered	Primarily nocturnal, hiding in soft substrate or submerged vegetation during the day.	Low	Watercourse on-site does not provide sufficient aquatic habitat to support species. No historical records for species in study area.
Bridle Shiner	Special Concern	Prefers clear water with abundant vegetation over silty or sandy vegetation	Low	Watercourse on-site does not provide sufficient aquatic habitat to support species. No historical records for species in study area.
Channel Darter	Special Concern	Prefers clear water with abundant vegetation over silty or sandy vegetation	Low	Watercourse on-site does not provide sufficient aquatic habitat to support species. No historical records for species in study area.

**TABLE C.7  
SCREENING RATIONALE FOR POTENTIAL SPECIES AT RISK ON-SITE OR WITHIN STUDY AREA**

Species	ESA Status	Habitat Use	Probability of Occurrence On-Site or Within Study Area	Rationale
Lake Sturgeon	Endangered	Large lakes and rivers. Forages in cool water, 4-9m deep over soft substrates. Spawns in shallower, fast-flowing areas over rocks or gravel.	Low	Watercourse on-site does not provide sufficient aquatic habitat to support species. No historical records for species in study area.
Northern Brook Lamprey	Special Concern	Prefers shallow areas with warm water. Larvae burrows in soft substrate for up to 7 years.	Low	Watercourse on-site does not provide sufficient aquatic habitat to support species. No historical records for species in study area.
River Redhorse	Special Concern	Prefers fast-flowing, clear rivers over rocky substrate	Low	Watercourse on-site does not provide sufficient aquatic habitat to support species. No historical records for species in study area.
Silver Lamprey	Special Concern	Larvae live 4-7 years in burrows, preference to soft substrate.	Low	Watercourse on-site does not provide sufficient aquatic habitat to support species. No historical records for species in study area.
<b>Insects</b>				
Bogbean Buckmoth	Endangered	Preferred food plant is bog bean, present in a variety of wetlands including bogs, swamps and fens.	Low	Preferred wetland habitat is not present on-site.
Gypsy Cuckoo Bumble Bee	Endangered	Inhabits a wide range of habitats: open meadows, agricultural and urban areas, boreal forests and woodlands.	Low	Currently the only known population is in Pinery Provincial Park.
Monarch Butterfly	Special Concern	Caterpillars require milkweed plants confined to meadow and open areas. Adult butterflies use more diverse habitat with a variety of wildflowers	Moderate	Potentially suitable foraging habitat available for Monarch on-site.
Mottled Duskywing	Endangered	Larval food plant (New Jersey Tea) found in sandy areas and alvars.	Low	Sandy areas and alvars not present in the study area.
Nine-spotted Lady Beetle	Endangered	Habitat generalist	Low	No recent occurrence reports in the area, thought to be locally extirpated
Rusty-patched Bumble Bee	Endangered	Habitat generalist	Low	Currently the only known population is in Pinery Provincial Park.
Traverse Lady Beetle	Endangered	Habitat generalist	Low	No new records of Traverse Lady Beetle in Ontario, species thought to be absent in former habitats.
West Virginia White Butterfly	Special Concern	Requires mature moist deciduous woods with larval host plant toothwort.	Low	Necessary vegetation and toothwort plant not present on-site or within study area.
Yellow-banded Bumble Bee	Special Concern	Habitat generalist; mixed woodlands, variety of open habitat	Moderate	Potentially suitable foraging habitat available for yellow-banded bumble bee on-site.



## **APPENDIX D**

Headwater Drainage Feature Assessment  
(GEMTEC, 2024)

May 10, 2024

File: 102151.001

TLC Holdings Ltd.  
1380 Howie Road  
Carp, Ontario  
K0A 1L0

Attention: Mr. Tom Carroll

**Re: Headwater Drainage Feature Assessment  
Proposed Plan of Subdivision, 3160 Carp Road  
City of Ottawa, Ontario**

---

## **INTRODUCTION**

GEMTEC Consulting Engineers and Scientists Limited (GEMTEC) was retained by TLC Holdings Ltd. to carry out a Headwater Drainage Feature Assessment (HDFA) for the property located at 3160 Carp Road, in the City of Ottawa, hereafter referred to as the “subject property”. This letter provides the methodologies and results of the HDFA conducted at the subject property. The subject property location and layout are illustrated on Figure A.1 and A.2, respectively, in Attachment A.

### **Purpose**

The proponent is seeking to develop a plan of subdivision on an approximately 23-hectare (ha) property located at 3160 Carp Road, in the City of Ottawa. The project involves the planned construction of 12 lots for future single-family residential dwellings, and a road.

### **Objective**

Under Section 28 (1) of the Conservation Authorities Act, conservation authorities have the ability to define the definition of a watercourse, which is defined under Section 28 (5) of the Act as “*An identifiable depression in the ground in which a flow of water regularly or continuously occurs*”. Headwater drainage features are defined as “*non-permanently flowing drainage features that may not have defined bed or banks; they are first-order and zero-order intermittent and ephemeral channels, swales and connected to headwater wetlands, but do not include rills or furrows*”. According to conservation authorities in Ontario, headwater drainage features (HDFs) meet the definition of a watercourse.

The objective of the work presented herein is twofold; 1) to identify HDFs and 2) to evaluate and classify any HDFs on-site, in accordance with “*Evaluation, Classification and Management of Headwater Drainage Features Guidelines*” from the Toronto Region Conservation Authority and

the Credit Valley Conservation (TRCA/CVC, 2014), and to recommend mitigation and conservation measures for HDFs present on-site.

## METHODOLOGY

### Desktop Review

A desktop information gathering exercise was completed to aid in the scoping of field surveys and to gather background information relating to HDFs on-site. Information relating to the presence and assessment of HDFs on-site was obtained from the following sources:

- Evaluation, Classification and Management of Headwater Drainage Features Guidelines (TRCA/CVC, 2014);
- Ontario Stream Assessment Protocol, Section 4, Module 11 (OSAP, 2017);
- Land Information Ontario (OMNR, 2011);
- Mississippi Valley Conservation Authority Interactive Map (MVCA, n.d.);
- City of Ottawa Official Plan (Ottawa, 2022); and
- Make a Map: Natural Heritage Areas (OMNRF, 2014).

### Field Investigations

Three field surveys were undertaken to evaluate the HDF identified on-site. Field surveys completed in support of this HDFA are outlined in Table 2.1 below. Photographs of site features taken during field investigations are provided in Attachment C.

**Table 2.1 Summary of Field Investigations**

Date	Time	Weather	HDFA Visit Number
April 27, 2023	12:45 – 16:00	12°C, ~75% cloud cover, Beaufort 1, no precipitation	1 (Spate)
May 17, 2023	11:00 – 13:00	6°C, ~25% cloud cover, Beaufort 3, no precipitation	2 (Spate)
July 26, 2023	07:15 – 08:45	19°C, ~25% cloud cover, Beaufort 1, no precipitation	3 (Baseflow)

### Headwater Drainage Feature Assessment

Field data collection of HDFs on-site followed the protocol outlined in Section 4: Module 11, “Unconstrained Headwater Sampling” from the Ontario Stream Assessment Protocol (OSAP) (Stanfield, 2017).

Data collected during the site investigations included flow conditions, sediment transport, feature roughness, riparian and feature vegetation, as well as upstream and downstream site features. As outlined in the OSAP manual for assessing HDFs, three site surveys were completed.

Classification of the HDFs on-site followed the protocols outlined in the Evaluation, Classification and Management of Headwater Drainage Features Guidelines manual (TRCA/CVC, 2014). Functions of the HDF that were evaluated included hydrology, vegetation, fish and fish habitat, and terrestrial habitat. Mitigation and management recommendations are provided for the HDFs based on the results of the classification.

## **HEADWATER DRAINAGE FEATURES ASSESSMENT**

### **Site Characteristics**

The 23 ha site currently consists of a mosaic of deciduous forest/woodland, cultural meadow, open agriculture, and thicket communities. The site is located within the 'Carp River' watershed and is under the jurisdiction of the Mississippi Valley Conservation Authority (MVCA).

Based on the desktop review and the site surveys, six HDFs occur on-site and are identified as HDF1, HDF2, HDF3, HDF3A, HDF3B and HDF4. All six HDFs are illustrated on Figure A.3 in Attachment A.

HDF1, HDF2, HDF3, and HDF3B and HDF4 are artifact of historical agricultural drainage efforts. HDF3A is associated with water pooling and groundwater drainage, likely due to snowmelt.

Each HDF is identified and described in more detail in the subsections below, while table B.1 in Attachment B provides a summary of collected field data.

### **HDF1**

HDF1 originates within the north of the property and flows along the property boundary in a northeasterly direction for approximately 52 m before exiting the subject property. From there, the feature continues off-site in a linear, northeastern direction along the agricultural fields for approximately 1.2 km.

HDF1 is a channelized feature that was observed to have standing water during the first two site investigations, and dry during the third. No evidence of seeps, springs or upwellings were noted during any of the three surveys. HDF1 had scrubland vegetation within the feature while the riparian zone was dominated by cropped fields.

Table B.1 in Attachment B summarizes the existing conditions and characteristics of HDF1 observed during the site surveys. During the site surveys, HDF1 was assessed as one continuous feature with no site break triggers.

### **HDF2**

HDF2 is an isolated feature that originates in the north of the property and flows in a southeast direction for approximately 50 m before abruptly ending.

HDF2 is a channelized feature that was observed to have standing water during the first assessment in April 2023, and dry during the second and third visits. No evidence of seeps, springs or upwellings were noted during any of the three surveys. HDF2 had scrubland vegetation within both the feature and riparian zone.

Table B.1 in Attachment B summarizes the existing conditions and characteristics of HDF2 observed during the site surveys. During the site surveys, HDF2 was assessed as one continuous feature with no site break triggers.

### **HDF3**

HDF3 originates in the eastern extent of the property and flows through the cultural thicket in a northwest direction for approximately 460 m before discharging into a vertical drain.

HDF3 is a channelized feature that was observed to have minimal flow during the first visit, interstitial flow during the second, and dry during the third. No evidence of seeps, springs or upwellings were noted during any of the three surveys. HDF3 had scrubland vegetation within both the feature and riparian zone.

Table B.1 in Attachment B summarizes the existing conditions and characteristics of HDF3 observed during the site surveys. During the site surveys, HDF3 was assessed in two segments based on site break triggers, but the segments displayed similar site features and conditions and have been grouped for evaluation purposes.

### **HDF3A**

HDF3A originates in cultural thicket in the northeast of the property and flows in a northern direction for approximately 45 m before discharging into the downstream extent of HDF3, and ultimately, the vertical drain.

HDF3A has no defined feature and was observed to have minimal flow during the first visit, interstitial flow during the second, and dry during the third. No evidence of seeps, springs or upwellings were noted during any of the three surveys. Vegetation within HDF3A and within the riparian zone was dominated by meadow.

Table B.1 in Attachment B summarizes the existing conditions and characteristics of HDF3A observed during the site surveys. During the site surveys, HDF3A was assessed as one continuous feature with no site break triggers.

### **HDF3B**

HDF3B is located within the eastern-central portion of the property and originates as an off-shoot from HDF3, within the cultural thicket. HDF3B continues from the confluence for approximately 20 m in a northeast direction before abruptly ending before approaching the agricultural fields.

HDF3B is a channelized feature that was observed to have standing water during the first two site investigations and was dry by the third. No evidence of seeps, springs or upwellings were noted during any of the three surveys. HDF3B had meadow vegetation within the feature while the riparian zone was dominated by scrubland.

Table B.1 in Attachment B summarizes the existing conditions and characteristics of HDF3B observed during the site surveys. During the site surveys, HDF3B was assessed as one continuous feature with no site break triggers.

#### **HDF4**

HDF4 originates in the southeastern corner of the property and flows in a southeastern direction for approximately 35 m before discharging into the permanent watercourse along the agricultural field.

HDF4 is a channelized feature that was observed to have interstitial flow during the first site investigation, standing water during the second, and was dry during the third. No evidence of seeps, springs or upwellings were noted during any of the three surveys. Vegetation within HDF4 and within the riparian zone was dominated by scrubland.

Table B.1 in Attachment B summarizes the existing conditions and characteristics of HDF4 observed during the site surveys. During the site surveys, HDF4 was assessed as one continuous feature with no site break triggers.

#### **CLASSIFICATION**

All HDFs on-site were classified based on the information collected during the site surveys pertaining to hydrology, riparian habitat, fish and fish habitat and terrestrial components. Using the linking classification to management flow chart provided by the TRCA and CVC (2014), illustrated in Figure 1 below, the classification of each HDF was used to determine management recommendations.



H3B and H4 were channelized and had standing water during the first and second visit, but dry conditions during third site investigation. Both features provide contributing fish habitat, limited terrestrial habitat, and important scrubland riparian habitat. As such **conservation** is required for the feature.

A summary of the classification and management recommendation for all HDFs is provided in Table B.2 in Attachment B.

## **MANAGEMENT RECOMMENDATIONS AND MITIGATION MEASURES**

In accordance with the guidance document (TRCA/CVC, 2014), HDFs classified as valued functions require **conservation**; these are typically features characterized by contributing hydrology, and important riparian vegetation. Based on the classification in Section 4 above, H3, H3B and H4 have been field verified to provide valued or contributing hydrology, contributing fish habitat, limited terrestrial habitat and important riparian habitat, as such **conservation** is required for these features.

As outlined in the guidance document, conservation management includes: maintaining, relocating, and/or enhancing the existing feature and riparian zone corridor; restoring lost functions through enhanced lot level controls; maintaining or replacing on-site flows using mitigation measures and/or wetland creation; maintaining or replacing external flows; and feature must remain connected to downstream features (TRCA/CVC, 2014).

In accordance with the guidance document (TRCA/CVC, 2014), HDFs classified as contributing functions require **mitigation**; these are typically features characterized by contributing hydrology, contributing fish habitat, contributing terrestrial habitat and/or contributing riparian habitat. Based on the classification in Section 4 above, H1 and H3A has been field verified to provide contributing hydrology and fish habitat, limited terrestrial habitat and/or valued riparian habitat, as such **mitigation** is required for this feature.

As outlined in the guidance document, mitigation management includes: replicating or enhancing feature functions through enhanced lot level conveyance measures (well vegetated swales), replicate on-site flow and outlet flows at the top of the system to maintain feature functions with vegetated swales/bioswales, if catchment drainage has been previously removed due to diversion of stormwater flows, restore lost functions through enhanced lot level controls, replicated functions by lot level conveyance measures (e.g. vegetated swales) connected to the natural heritage system, as feasible and/or Low Impact Development (LID) stormwater options (TRCA/CVC, 2014).

In accordance with the guidance document (TRCA/CVC, 2014) HDFs classified as limited functions require **no management**; these are typically features with no or minimal flow and/or no fish or fish habitat. Based on the classification in Section 4 above, H2 has been field verified to

have very limited hydrology, fish habitat, terrestrial habitat and/or riparian habitat, as such **no management** is required for this feature.

In addition to the management recommendations outlined above, the following mitigation measures are provided by GEMTEC in order to minimize or eliminate potential impacts to fish habitat.

- All future development and construction activities within the study area, including ditching, culvert installation, erosion and sediment control and storm water management should be completed in accordance with Ontario Provincial Standard Specification 182 and OPSS 805.
- No in-water work should occur between March 15 and June 30 of any year to protect spawning fish habitat adjacent to the development area. All in-water habitat features, including aquatic vegetation, natural woody debris and boulders should be left in their current locations in the near shore area.
- When native soil is exposed, sediment and erosion control work in the form of heavy-duty sediment fencing shall be positioned along the down gradient edge of any construction envelopes adjacent to waterbodies.
- The development plan should include lot-side swales and/or road side ditches designed to promote infiltration.
- In order to protect fish habitat from contamination, it is recommended that all machinery be maintained in good working condition and that all machinery be fueled a minimum of 30 m from the high water mark.
- Any temporary storage of aggregate material shall be set back from the water's edge by no less than 40 m and be contained by heavy-duty silt fencing.

## SUMMARY

A headwater drainage feature assessment was completed and six HDFs were identified on-site, identified as HDF1, HDF2, HDF3, HDF3A, HDF3B, and HDF4. Conservation is required for H3, HDF3B, and HDF4. Conservation management should include: maintaining, relocating and/or enhancing the existing features or riparian zone corridor; restoring lost functions through enhanced lot level controls; maintaining or replacing on-site flows through mitigation; maintaining or replacing external flows and maintaining connectivity with downstream features. Mitigation management is required for HDF1 and H3A. Mitigation management should include: replicating or enhancing feature functions through enhanced lot level conveyance measures. No management is required for HDF2 .

We trust this report provides sufficient information for your present purposes. If you have any questions concerning this report, please do not hesitate to contact the undersigned.

Sincerely,

Handwritten signature of Emily Pentz in blue ink.

Emily Pentz, B.Sc.  
Junior Biologist

Handwritten signature of Taylor Warrington in black ink.

Taylor Warrington, B.Sc.  
Biologist

## REFERENCES

Ontario (Ontario). 1990. Conservation Authorities Act. R.S.O. 1990. Chapter C.27. Last amendment: 2019, c.9. Sched 2.

Ontario Ministry of Natural Resources (OMNR). 2011. Land Information Ontario (LIO).

Ontario Ministry of Natural Resources and Forestry (OMNRF). 2014. Make a Map: Natural Heritage Areas.

Mississippi Valley Conservation Authority (MVCA). n.d. Interactive Property Map, viewed online March 6, 2024. Available: <https://mvc.on.ca/development-permits/interactive-property-map/>

Stanfield, L (Ed.). 2017. Ontario Stream Assessment Protocol (OSAP). Version 10. Ontario, Canada.

Ottawa, City of (Ottawa). 2022. City of Ottawa Official Plan.

Toronto Region Conservation Authority and Credit Valley Conservation (TRCA/CVC). 2014. Evaluation, Classification and management of Headwater Drainage Features Guidelines. TRCA Approval July 2013, Finalized January 2014.



## **ATTACHMENT A**

Figure A.1 – Site Location



Figure A.2 – Site Layout

Figure A.3 – Headwater Drainage Feature Assessment



**Legend**

- Property Boundary
- Study Area



**GEMTEC**  
CONSULTING ENGINEERS  
AND SCIENTISTS

32 Steacie Drive,  
Ottawa, ON K2K 2A9  
T: (613) 836-1422  
www.gemtec.ca  
ottawa@gemtec.ca

Client: <b>TLC Holdings Ltd.</b>	Project: <b>102151.001</b>
----------------------------------	----------------------------

Location  
**3160 Carp Road  
Ottawa, Ontario**

Drwn By: EP	Chkd By: TW	<b>Site Location</b>
----------------	----------------	----------------------


Date: April 2024	Rev. 0	<b>Figure: A.1</b>
© Queen's Printer for Ontario		

Coordinate System: NAD 1983 UTM Zone 18N  
 Service Layer Credits: World Street Map: City of Ottawa, Province of Ontario, Esri Canada, Esri, TomTom, Garmin, SafeGraph, GeoTechnologies, Inc, METI/NASA, USGS, EPA, NPS, US Census Bureau, USDA, NRCAN, Parks Canada  
 World Topographic Map: City of Ottawa, Province of Ontario, Esri Canada, Esri, TomTom, Garmin, SafeGraph, METI/NASA, USGS, EPA, NPS, USDA, NRCAN, Parks Canada



### Legend

- Property Boundary
- Study Area
- Watercourse
- Headwater Drainage Feature
- Development Concept



**GEMTEC**  
CONSULTING ENGINEERS  
AND SCIENTISTS

32 Steacie Drive,  
Ottawa, ON K2K 2A9  
T: (613) 836-1422  
www.gemtec.ca  
ottawa@gemtec.ca

Client: <b>TLC Holdings Ltd.</b>	Project: 102151.001
----------------------------------	---------------------

Location  
**3160 Carp Road  
Ottawa, Ontario**

Drwn By: EP	Chkd By: TW	<b>Site Layout</b>
----------------	----------------	--------------------

Date: April 2024	Rev. 0	<b>Figure: A.2</b>
© Queen's Printer for Ontario		

Coordinate System: NAD 1983 UTM Zone 18N  
 Service Layer Credits: Hybrid Reference Layer: Esri Community Maps Contributors, City of Ottawa, Province of Ontario, Esri Canada, Esri, TomTom, Garmin, SafeGraph, GeoTechnologies, Inc, METI/NASA, USGS, EPA, NPS, US Census Bureau, USDA, NRCAN, Parks Canada  
 World Imagery: Maxar




**Legend**

- Property Boundary
- Study Area
- Watercourse
- Development Concept

**Headwater Drainage Features**

- Conservation
- Mitigation
- No Management Required



32 Steacie Drive,  
Ottawa, ON K2K 2A9  
T: (613) 836-1422  
www.gemtec.ca  
ottawa@gemtec.ca

Client:	TLC Holdings Ltd.	Project:	102151.001
---------	-------------------	----------	------------

Location  
**3160 Carp Road  
Ottawa, Ontario**

Drwn By: EP	Chkd By: TW	<b>Headwater Drainage Feature Assessment</b>
----------------	----------------	--

Date: April 2024	Rev. 0	<b>Figure: A.3</b>
------------------	-----------	--------------------

Coordinate System: NAD 1983 UTM Zone 18N  
 Service Layer Credits: World Imagery: Maxar, Microsoft  
 Hybrid Reference Layer: Esri Community Maps Contributors, City of Ottawa, Province of Ontario, Esri Canada, Esri, TomTom, Garmin, SafeGraph, GeoTechnologies, Inc, METI/NASA, USGS, EPA, NPS, US Census Bureau, USDA, NRCAN, Parks Canada



## **ATTACHMENT B**

Headwater Drainage Feature Assessment Summary Tables

**Table B.1  
Summary of Headwater Drainage Features**

Site Visit	Hydrology			Vegetation Assessment		Channel Form			Sediment Transport		
	Flow Influence (FI)	Flow Condition (FC)	Feature Type (FT)	Feature	Riparian	Average Wetted Width (m)	Average Depth (range) (cm)	Average Bankfull Width (m)	Substrate	Sediment Transport	Sediment Deposition
<b>HDF1</b>											
1	Freshet (1)	Standing Water (2)	Channelized (2)	Scrubland (5)	Cropped (3)	1.25	12	1.6	Sandy Silt	None	None
2	Spate (2)	Standing Water (2)	Channelized (2)	Scrubland (5)	Cropped (3)	0.8	7	1.6	Sandy Silt	None	None
3	Baseflow (3)	Dry (1)	Channelized (2)	Scrubland (5)	Cropped (3)	--	--	1.6	Sandy Silt	None	None
<b>HDF2</b>											
1	Freshet (1)	Standing Water (2)	Channelized (2)	Scrubland (5)	Scrubland (5)	1	7	1.3	Sandy Silt	None	None
2	Spate (2)	Dry (1)	Channelized (2)	Scrubland (5)	Scrubland (5)	--	--	1.3	Sandy Silt	None	None
3	Baseflow (3)	Dry (1)	Channelized (2)	Scrubland (5)	Scrubland (5)	--	--	1.3	Sandy Silt	None	None
<b>HDF3</b>											
1	Freshet (1)	Minimal Flow (4)	Channelized (2)	Scrubland (5)	Scrubland (5)	1.5	14	2	Sandy Silt	None	None
2	Spate (2)	Interstitial Flow (3)	Channelized (2)	Scrubland (5)	Scrubland (5)	1.75	10.5	2	Sandy Silt	None	None
3	Baseflow (3)	Dry (1)	Channelized (2)	Scrubland (5)	Scrubland (5)	--	--	2	Sandy Silt	None	None
<b>HDF3A</b>											
1	Freshet (1)	Minimal Flow (4)	No Defined Feature (4)	Meadow (4)	Meadow (4)	1	14	--	Sandy Silt	None	None
2	Spate (2)	Interstitial Flow (3)	No Defined Feature (4)	Meadow (4)	Meadow (4)	0.9	12	--	Sandy Silt	None	None
3	Baseflow (3)	Dry (1)	No Defined Feature (4)	Meadow (4)	Meadow (4)	--	--	--	Sandy Silt	None	None
<b>HDF3B</b>											
1	Freshet (1)	Standing Water (2)	Channelized (2)	Meadow (4)	Scrubland (5)	1.25	10	1.25	Sandy Silt	None	None
2	Spate (2)	Standing Water (2)	Channelized (2)	Meadow (4)	Scrubland (5)	1.25	8.5	1.25	Sandy Silt	None	None
3	Baseflow (3)	Dry (1)	Channelized (2)	Meadow (4)	Scrubland (5)	--	--	1.25	Sandy Silt	None	None
<b>HDF4</b>											
1	Freshet (1)	Interstitial Flow (3)	Channelized (2)	Scrubland (5)	Scrubland (5)	0.8	8	1	Sandy Silt	None	None
2	Spate (2)	Standing Water (2)	Channelized (2)	Scrubland (5)	Scrubland (5)	0.8	6	1	Sandy Silt	None	None
3	Baseflow (3)	Dry (1)	Channelized (2)	Scrubland (5)	Scrubland (5)	--	--	1	Sandy Silt	None	None

**Table B.2**  
**Summary of HDF Classification and Management Recommendations**

HDF	Step 1		Step 2	Step 3	Step 4	Management Recommendation
	Hydrology	Modifiers	Fish Habitat	Terrestrial Habitat	Riparian Vegetation	
H1	Contributing- Ephemeral	None	Contributing	Limited	Limited -Cropped	Mitigation
H2	Limited	None	Contributing	Limited	Important - Scrubland	No Management Required
H3	Valued - Intermittent	None	Contributing	Limited	Important - Scrubland	Conservation
H3A	Valued - Intermittent	None	Contributing	Limited	Valued - Meadow	Mitigation
H3B	Contributing- Ephemeral	None	Contributing	Limited	Important - Scrubland	Conservation
HDF4	Contributing- Ephemeral	None	Contributing	Limited	Important - Scrubland	Conservation




**ATTACHMENT C**

Photolog



Site Photograph 1: Mixed Meadow (MEM).



Site Photograph 2: Fresh – Moist Lowland Deciduous Forest (FODM7).



Site Photograph 3: Open Agriculture (OAG).



Site Photograph 4: Meadow (ME).



Site Photograph 5: Upstream extent of HDF3 and Buckthorn Deciduous Shrub Thicket (THDM2-6).



Site Photograph 6: The vertical drain on-site.



Site Photograph 7: HDF1



Site Photograph 8: HDF2



Site Photograph 9: Downstream extent of HDF3.



Site Photograph 10: HDF3A



Site Photograph 11: HDF3B



Site Photograph 12: HDF4



## **APPENDIX E**

MNRF General Habitat Descriptions

i. Blanding's Turtle

# General Habitat Description for the Blanding's Turtle (*Emydoidea blandingii*)

A general habitat description is a technical document that provides greater clarity on the area of habitat protected for a species based on the general habitat definition found in the Endangered Species Act, 2007. General habitat protection does not include an area where the species formerly occurred or has the potential to be reintroduced unless existing members of the species depend on that area to carry out their life processes. A general habitat description also indicates how the species' habitat has been categorized, as per the policy "Categorizing and Protecting Habitat Under the Endangered Species Act", and is based on the best scientific information available.

## HABITAT CATEGORIZATION

- |   |   |
|---|---|
| 1 | Nest and the area within 30 m or Overwintering sites and the area within 30 m   |
| 2 | The wetland complex (i.e. all suitable wetlands or waterbodies within 500 m of each other) that extends up to 2 km from an occurrence, and the area within 30 m around those suitable wetlands or waterbodies |
| 3 | Area between 30 m and 250 m around suitable wetlands/waterbodies identified in Category 2, within 2 km of an occurrence   |

### Category 1

Nest sites and overwintering sites are essential features and along with the 30 m area surrounding them are considered to have the lowest tolerance to alteration. Blanding's Turtles depend on these areas for sensitive life processes including egg-laying, incubation, hatching of young, and hibernation. A 30 m radius (average tree height) buffer around nesting and overwintering sites is important to maintain the microclimate conditions (e.g., thermal, vegetative and lighting features). These areas are habitually used and may support concentrations of individuals.

#### *Nesting Sites*

Blanding's Turtle nests are created in open habitats with low vegetation cover and high sun exposure such as in forest clearings, meadows, shorelines, beaches, rock outcrops, cornfields, gravel roads, road shoulders, ploughed fields, gardens, powerline rights-of-ways, yards and abandoned railroad beds ( Linck et al. 1989, Ross and Anderson 1990, Kiviat 1997, Standing et al. 1999, Joyal et al. 2001, Congdon et al. 2008, Downing et al. 2010, Refsnider and Linck 2012). Females often show high fidelity to the same general nesting areas (Congdon et al. 1983, McNeil 2002, Congdon et al. 2011).

### *Overwintering Sites*

Overwintering sites are typically occupied for at least six months during the overwintering period in Ontario (Edge *et al.* 2009, Edge *et al.* 2010, Davy 2011 unpublished data, Paterson unpublished data 2013, NHIC 2013). Blanding's Turtles display overwintering site fidelity, using some sites year after year (Power 1989, McNeil 2002, Caverhill 2006 in Newton and Herman 2009, Edge *et al.* 2009). Many individuals may aggregate at one site while overwintering (Anderson 1990, St-Hilaire 2003 in COSEWIC 2005, Ross and Congdon *et al.* 2008, Edge *et al.* 2009).

Suitable Blanding's Turtle overwintering habitat typically includes permanent bogs, fens, marshes, ponds, channels or other habitats with free (unfrozen) shallow water (Joyal *et al.* 2001, Edge 2010, Seburn 2010). Blanding's Turtles studied in Algonquin Provincial park overwintered in wetlands with free water depths of 7 cm - 50 cm (Edge *et al.* 2009). This species may also hibernate within graminoid shallow marsh areas of larger marsh complexes by burying into substrates in areas of pooled water (Gillingwater unpublished data 2013). Blanding's Turtle's may also overwinter in seasonal pools or small excavated areas with standing water (Joyal *et al.* 2001, Rouse unpublished data 2012).

### **Category 2**

The wetland complex that extends up to 2 km from an occurrence and 30 m around these suitable wetlands/waterbodies (Category 2) will be considered to have a moderate level of tolerance to alteration before their function is compromised. For the purpose of general habitat protection for Blanding's Turtle, a wetland complex is defined as all wetlands that are within 500 m of each other. This definition is based on the biology of the species and its documents movement patterns between adjacent suitable wetlands/waterbodies. In cases where an occurrence is not within suitable aquatic habitat, the nearest wetland should be considered the starting point for delineating the wetland complex.

Blanding's Turtles depend on these wetlands and the surrounding habitat throughout their home range for life processes including feeding, mating, thermoregulation, movement, and protection from predators.

Blanding's Turtle home range sizes and lengths in Ontario vary significantly between individuals within the same population and between different populations. In Algonquin Provincial Park, the average range length of radio-tracked Blanding's Turtles was 1.8 km (1.2 standard deviation), with a maximum of 4.3 km (Edge 2013 unpublished data). Recent Ontario studies documented a 90th percentile home range length of radio-tracked Blanding's Turtles in Parry Sound District and Bancroft District of 2.0 and 2.3 km, respectively (Rouse unpublished data 2013, Cameron unpublished data 2013). Average range length of a population on Grenadier Island, Ontario, was 813 m, with a maximum range length just over 2 km. In a Minnesota population, average range length was just over 1.6 km, with a maximum range length just over 5 km (Pappas *et al.* 2000).

Blanding's Turtles regularly move between wetlands or other aquatic areas in order to access mates, overwintering sites, nesting sites, other seasonally required resources and thermoregulation sites (Congdon *et al.* 2008, Edge *et al.* 2010). In a study from Algonquin Provincial Park, Blanding's Turtles made an average of four movements between wetlands each year with an average movement distance of 231 m for males and 497 m for females (Edge *et al.* 2010). Average interwetland movement distances of a population in Maine was  $680 \pm 550$  m (Joyal *et al.* 2001). Rouse and Cameron (unpublished data 2013) found that Blanding's Turtles primarily moved through wetlands and other water and were rarely located more than 200 m from water. Since interwetland movements tend to average about 500 m, wetlands that are separated by more than 500 m from other suitable wetlands have a lower likelihood of being occupied.

A 30 m radius (average tree height) buffer around suitable wetlands helps to maintain microclimate conditions. Buffers of 30 m are widely recognized as providing a range of functional benefits to aquatic features and wetlands such as maintaining water quality by filtering sediment and nutrients, input of woody debris, and cooling water temperatures by shading and infiltrating surface runoff (OMNR 2010). Blanding's Turtles have also been shown to generally bask within 30 m of wetlands (Joyal *et al.* 2001).

Suitable habitat for Blanding's Turtles during the active season includes a variety of wetlands such as marsh, swamps, ponds, fens, bogs, slow-flowing streams, shallow bays of lakes or rivers, as well as graminoid shallow marsh and slough forest habitats that are adjacent to larger marsh complexes (Joyal *et al.* 2001, Gillingwater 2001, Gillingwater and Piraino 2004, 2007, Congdon *et al.* 2008, Edge *et al.* 2010; Seburn 2010). Suitable wetlands used during the active season are typically eutrophic (mineral or organic nutrient-rich), shallow with a soft substrate composed of decomposing materials, and often have emergent vegetation, such as water lilies and cattails (COSEWIC 2005, Congdon *et al.* 2008).

### **Category 3**

The area between 30 m and 250 m around suitable Category 2 wetlands/waterbodies will be considered to have the highest tolerance to alteration. Blanding's Turtles depend on these areas as movement corridors between wetlands, which are essential for carrying out life processes associated with Category 1 and 2 habitats.

Blanding's Turtle nests are typically close to permanent wetlands and reported average distances between nests and the nearest wetland range from 99.5 to 242 m, with maximum distances of 256 m to just over 400 m (Joyal *et al.* 2001, Beaudry *et al.* 2010, Congdon *et al.* 2011, Paterson *et al.* 2012, Refsnider and Linck 2012). Consequently, the area within 250 m of suitable aquatic habitat provides critical movement corridors through which hatchling Blanding's Turtles access wetlands after hatching. This habitat is also used by some hatchlings as overwintering habitat in their first year (Paterson *et al.* 2012).

Although Blanding's Turtles nest close to water, they often travel considerable distances from their wetland of origin during nesting migrations, with movements of 6 km being documented in some Ontario populations (Edge *et al.* 2010). Although wetlands and ponds are used as movement corridors when available, females make extensive movements through upland habitat to access nesting sites (Congdon *et al.* 2008). As mentioned in the previous section (see Category 2), Blanding's Turtles also make regular overland movements between wetlands throughout the active season in order to access Category 1 and 2 habitats within their home range. Category 3 habitat provides essential movement corridors of up to 500 m between wetlands, which will encompass the areas that are most likely to be used for overland movement.

## Activities in Blanding's Turtle habitat

Activities in general habitat can continue as long as the *function of these areas for the species is maintained and individuals of the species are not killed, harmed, or harassed.*

### Generally compatible:

- Recreational use of the water such as swimming, boating, and fishing.
- Small-scale alterations to land cover that do not impede overland movements or impair nesting sites.

### Generally not compatible\*:


- Significant draining, infilling, dredging, or other significant alteration of wetlands or other suitable waterbodies.
- Significant alteration of shorelines, especially hardening (e.g. the use of gabion baskets, rip-rap, and rock armour).

\* If you are considering an activity that may not be compatible with general habitat, please contact your local MNR office for more information.

## Key terms:

- **Thermoregulation:** Some animals, such as turtles, use thermoregulation to alter their internal body temperature through behavioural patterns, such as basking in the sun to increase body temperature or seeking out cool areas to lower body temperature.

### Sample application of the general habitat protection for Blanding's Turtle




## References

- Beaudry, F., P.G. DeMaynadier and M.L. Hunter Jr. 2010. Nesting movements and the use of anthropogenic nesting sites by Spotted Turtle (*Clemmys guttata*) and Blanding's Turtle (*Emydoidea blandingii*). *Herpetological Conservation and Biology* 5 (1): 1-8
- Cameron, G. 2013. Unpublished data. Species at Risk Biologist, Ontario Ministry of Natural Resources.
- Caverhill, B.P. 2006. Blanding's turtle conservation in Nova Scotia: linking science and stewardship through public education. M.Sc. thesis, Department of Biology, Acadia University, Wolfville, N.S.
- Congdon, J.D., D.W. Twinkle, G.L. Breitenbach and R.C. Van Loben Sels. 1983. Nesting ecology and hatching success in the turtle *Emydoidea blandingii*. *Herpetologica* 39(4):417-429.
- Congdon, J.D., Graham, T.E., Herman, T.B., Lang, J.W., Pappas, M.J., and Brecke, B.J. 2008. *Emydoidea blandingii* (Holbrook 1838) – Blanding's Turtle. In: Rhodin, A.G.J., Pritchard, P.C.H., van Dijk, P.P., Saumure, R.A., Buhlmann, K.A., and Iverson, J.B. (Eds.). *Conservation Biology of Freshwater Turtles and Tortoises: A Compilation Project of the IUCN/SSC Tortoise and Freshwater Turtle Specialist Group*. Chelonian Research Monographs No. 5, pp. 015.1-015.12, doi:10.3854/crm.5.015. blandingii.v12008, <http://www.iucn-tftsg.org/cbftt/>.
- Congdon, J.D., O.M. Kinney and R.D. Nagle. 2011. Spatial ecology and core-area protection of Blanding's Turtle (*Emydoidea blandingii*). *Canadian Journal of Zoology* 89: 1098-1106
- COSEWIC 2005. COSEWIC assessment and update status report on the Blanding's Turtle *Emydoidea blandingii* in Canada. Committee on the Status of Endangered Wildlife in Canada. Ottawa. 32 pp.
- Davy, C. 2011. Unpublished data from survey and monitoring work at Rondeau Provincial Park. University of Toronto, Ontario.
- Dowling, Z. Hartwig, T. Kiviat, E. and Keesing, F. 2010. Experimental Management of Nesting Habitat for Blanding's Turtle (*Emydoidea blandingii*). *Ecological Restoration* 28:2.
- Edge, C.B., Steinberg, B.D., Brooks, R.J., and Litzgus, J.D. 2010. Habitat Selection by Blanding's Turtles (*Emydoidea blandingii*) in relatively pristine landscape. *Ecoscience* 17(1):90-99.
- Edge, C.B., Steinberg, B.D., Brooks, R.J., and Litzgus, J.D. 2009. Temperature and site selection by Blanding's Turtles (*Emydoidea blandingii*) during hibernation near the species northern range. *Canadian Journal of Zoology* 87:825-834.
- Edge, C.B. 2013 Unpublished data from MSc. Research in Algonquin Provincial Park, Ontario.
- Gillingwater, S. D. 2001. A Selective Herpetofaunal Survey Inventory and Biological Research Study of Rondeau Provincial Park. Report submitted to ESRF, World Wildlife Fund. 94 pp.
- Gillingwater, S.D. 2013. Unpublished data from long-term survey and monitoring work provided through e-mail correspondence to Joe Crowley. Species at Risk Biologist, Upper Thames River Conservation Authority.

- Gillingwater, S.D. and T.J. Piraino. 2004. Chelonian Survey and Research Study of the Big Creek National Wildlife Area (2003) and Selective Herpetofaunal Survey, Inventory and Research Study of the Long Point National Wildlife Area (1996-1999, 2003). Final report submitted to the Canadian Wildlife Service. 65+pp.
- Gillingwater, SD and TJ Piraino. 2007. Turtle Research and Herpetofaunal Survey of the Long Point National Wildlife Area Update Report 2007. Report submitted to Canadian Wildlife Service
- Joyal, L.A., M. McCollough and M.L. Hunter Jr. 2001. Landscape ecology approaches to wetland species conservation: a case study of two turtle species in southern Maine. *Conservation Biology* 15(6): 1755-1762
- Kiviat, E., G. Stevens, R. Brauman, S. Hoeger, P.J. Petokas and G.G. Hollands. 2000. Restoration of wetland and upland habitat for the Blanding's turtle, *Emydoidea blandingii*. *Chelonian Conservation Biology* 3:650-657.
- Natural Heritage Information Centre (NHIC). 2013. Biodiversity Explorer: Species Lists, Element Occurrence and Natural Areas databases. Natural Heritage Information Centre, Ontario Ministry of Natural Resources, Peterborough, Ontario.
- Newton, E.J. and Herman, T.B. 2009. Habitat, movements, and behaviour of overwintering Blanding's Turtles (*Emydoidea blandingii*) in Nova Scotia. *Canadian Journal of Zoology* 87:299-309.
- OMNR. 2010. Forest Management Guide for Conserving Biodiversity at the Stand and Site Scales. Toronto: Queen's Printer for Ontario. 211 pp.
- Paterson, J. 2013. Unpublished data from M.Sc. research. Laurentian University, Sudbury, Ontario.
- Paterson, J.E., B.D. Steinberg and J.D. Litzgus. 2012. Revealing a cryptic life-history stage: difference in habitat selection and survivorship between hatchlings of two turtle species at risk (*Glyptemys insculpta* and *Emydoidea blandingii*).
- Power, T. 1989. Seasonal movements and nesting ecology of a relict population of Blanding's turtle (*Emydoidea blandingii* (Holbrook)) in Nova Scotia. M.Sc. thesis, Department of Biology, Acadia University, Wolfville, N.S.
- Piepgas, S. A., and J. W. Lang. 2000. Spatial ecology of Blanding's Turtle in central Minnesota. *Chelonian Conservation and Biology* 3(4):589-601.
- Refsnider, J.M. and M. H. Linck. 2012. Habitat use and movement patterns of Blanding's Turtles (*Emydoidea blandingii*) in Minnesota, USA: a landscape approach to species conservation. *Herpetological Conservation and Biology* 7(2): 185-192.
- Ross, D.A. and R.K. Anderson. 1990. Habitat use, movements, and nesting of *Emydoidea blandingii* in central Wisconsin. *Journal of Herpetology* 24:6-12.
- Rouse, J. 2013. Unpublished data. Species at Risk Biologist, Ontario Ministry of Natural Resources.
- Seburn, D.C. 2010. Blanding's Turtle, *Emydoidea blandingii*, Habitat Use During Hibernation in Eastern Ontario. *The Canadian Field-Naturalist* 124(3): 263-265.
- Standing, K.L., T.B. Herman and I.P. Morrison. 1999. Nesting ecology of Blanding's turtle (*Emydoidea blandingii*) in Nova Scotia, the northeastern limit of the specie's range. *Canadian Journal of Zoology* 77:1609-1614.

experience • knowledge • integrity



civil  
geotechnical  
environmental  
field services  
materials testing

civil  
géotechnique  
environnementale  
surveillance de chantier  
service de laboratoire des matériaux

expérience • connaissance • intégrité

