

Cardinal Creek Village South Existing Conditions Report

Final Report

November 5, 2024

Submitted To:

Mike Green
Taggart (Tamarack Developments)
3187 Albion Road South
Ottawa, ON
K1V 8Y3

KILGOUR & ASSOCIATES LTD.
www.kilgourassociates.com



TABLE OF CONTENTS

1.0 INTRODUCTION	1
2.0 ENVIRONMENTAL POLICY CONTEXT	1
2.1 THE PROVINCIAL POLICY STATEMENT, 2020, 2024	1
2.2 THE CITY OF OTTAWA OFFICIAL PLAN (2021)	3
2.3 CONSERVATION AUTHORITIES ACT, 1990	4
2.4 SPECIES AT RISK ACT, 2002	4
2.5 ENDANGERED SPECIES ACT, 2007	4
2.6 FISHERIES ACT, 1985	5
2.7 MIGRATORY BIRDS CONVENTION ACT, 1994	5
2.8 FISH AND WILDLIFE CONSERVATION ACT, 1997	5
2.9 GREATER CARDINAL CREEK SUBWATERSHED MANAGEMENT PLAN	5
3.0 PROPERTY IDENTIFICATION	6
4.0 METHODOLOGY.....	7
4.1 DESKTOP AND BACKGROUND DATA REVIEW	7
4.1.1 General Records Review.....	7
4.1.2 Species at Risk Screening.....	7
4.1.3 Site-Specific Background Data Review	8
4.1.4 Agency Consultation	9
4.2 FIELD SURVEYS	9
4.2.1 Site Work Summary	9
4.2.2 Surface Water Characterization.....	11
4.2.3 Ecological Land Classification	11
4.2.4 Tree Inventory, Butternut and Black Ash	11
4.2.5 Breeding Birds.....	12
4.2.6 Bats and Other Mammals.....	12
5.0 RESULTS	13
5.1 GENERAL NATURAL HERITAGE CONTEXT	13
5.2 LANDFORMS, SOILS, AND GEOLOGY	13
5.3 SURFACE WATER, GROUNDWATER AND FISH HABITAT	13
5.3.1 General Context	13
5.3.2 Headwater Drainage Features.....	15
5.4 ECOLOGICAL LAND CLASSIFICATION	19
5.4.1 Open Agriculture – Annual Row Crops (OAGM1)	20
5.4.2 Fresh – Moist Sugar Maple – Hardwood Deciduous Forest Type (FODM6-5)	20
5.4.3 Dry – Fresh Deciduous Woodland Ecosite (WODM4)	22
5.4.4 Reed Canary Grass Graminoid Meadow Type (MEGM3-8).....	24
5.4.5 Native Deciduous Regeneration Thicket Type (THDM4-1)	26
5.4.6 Dry – Fresh Sugar Maple – Hardwood Deciduous Forest Type (FODM5-9)	27
5.4.7 Cultural Meadow Ecosite (CUM1-1)	29
5.4.8 Constructed (CV).....	30



5.5 WILDLIFE SURVEYS	31
5.5.1 Breeding Birds.....	31
5.5.2 Bats and Other Mammals.....	32
5.6 SPECIES AT RISK	33
5.6.1 SAR Bats.....	35
5.6.2 Black Ash.....	35
5.7 SIGNIFICANT NATURAL HERITAGE FEATURES	36
5.7.1 Significant Woodlands and Canopy Cover.....	36
5.7.2 Significant Valleylands.....	39
5.8 SIGNIFICANT WILDLIFE HABITAT	39
5.8.1 Seasonal Concentration Areas.....	39
5.8.2 Rare Vegetation Communities or Specialized Habitat for Wildlife.....	39
5.9 OTHER NATURAL HERITAGE FEATURES	40

6.0 OPPORTUNITIES.....	40
7.0 CLOSURE.....	42
8.0 LITERATURE CITED.....	43
LITERATURE CITED.....	10

List of Figures

Figure 1 Site Location.....	2
Figure 2 Field Study Locations.....	10
Figure 3 Existing Conditions.....	14
Figure 4 Fresh – Moist Sugar Maple – Hardwood Deciduous Forest Type (FODM6-5).....	21
Figure 5 FODM6-5 Soil Core.....	22
Figure 6 Dry – Fresh Woodland Ecosite (WODM4).....	23
Figure 7 Soil Core in WODM4.....	24
Figure 8 Reed Canary Grass Graminoid Meadow Type (MEGM3-8).....	25
Figure 9 MEGM3-8 Soil Core.....	26
Figure 10 Native Deciduous Regeneration Thicket Type (THDM4-1).....	27
Figure 11 Dry – Fresh Sugar Maple – Hardwood Deciduous Forest Type (FODM5-9).....	28
Figure 12 FODM5-9 Soil Core.....	29
Figure 13 Cultural Meadow (CUM1-1) adjacent to Open Agriculture (OAGM1-1).....	30
Figure 14 Constructed (CV).....	31
Figure 15 Significant Natural Heritage Features.....	37

List of Tables

Table 1 Field Study Dates.....	9
Table 2 Hydrology Classifications for HDFs.....	17
Table 3 Riparian Classifications.....	18



Table 4 Fish and Fish Habitat Classification.....	18
Table 5 Terrestrial Classifications	19
Table 6 Breeding Bird Survey Data.....	32
Table 7 Acoustic Bat Survey Data.....	33
Table 8 Species at risk with moderate or high potential to interact with the project	34
Table 9 Characterization of Significant Woodland Areas.....	36
Table 10 Assessment of Canopy Tree Benefits on the Site.....	38

List of Appendices

Appendix A Qualifications of Report Authors
Appendix B Species at Risk Screening

List of Acronyms and Abbreviations

cm – centimetres
e.g. – *exempli gratia*
ECR – Existing Conditions Report
EIS – Environmental Impact Study
ELC – Ecological Land Classification
ESA – *Endangered Species Act*
FWCA – *Fish and Wildlife Conservation Act*
ha – hectare
i.e. – id est
KAL – Kilgour & Associates Ltd.
km – kilometre
m – metre
MBCA – *Migratory Birds Convention Act*
MECP – Ministry of Environment, Conservation and Parks
MNR – Ministry of Natural Resources and Forestry
NHIC – Natural Heritage Information Centre
PPS – Provincial Policy Statement
SAR – species at risk
SARA – *Species at Risk Act*
SWH – Significant Wildlife Habitat



1.0 INTRODUCTION

This report is an Existing Conditions Report (ECR) prepared by Kilgour & Associates Ltd. (KAL; Appendix A) on behalf of Taggart (Tamarack Developments; “the Client”) in support of a future residential development known as Cardinal Creek Village South (the “Site”; Figure 1). The Site is located at 1296 and 1400 Old Montreal Road and encompasses the northern portion of each property, from Cardinal Creek northward.

Natural heritage review work in support of development planning began in 2011, with several minor updates taking place over the subsequent years. An Environmental Impact Study (EIS) was prepared in 2021 by Muncaster Environmental Planning Inc.; however, the City of Ottawa updated its EIS guidelines in 2023 and an update is anticipated for development approval. This ECR will provide a basis for a subsequent EIS. This ECR will also serve to identify opportunities for consideration in the planning process related to development options for the Site.

Accordingly, this report identifies natural heritage conditions on the Site based on field studies performed to date and reviews of publicly available records and data for the area. The report also outlines the policy context associated with future development plans. The content of this report (i.e. the natural heritage system review) was completed per the *Environmental Impact Study Guidelines* (City of Ottawa, 2023).

A future EIS based on this report will be required to:

- Identify natural heritage features on or adjacent to the Site;
- Assess potential impacts of the proposed development to existing features; and
- Recommend mitigation measures to minimize or eliminate identified impacts.

This current ECR addresses the first component.

2.0 ENVIRONMENTAL POLICY CONTEXT

Natural heritage policies and legislation relevant to the development of the Site are outlined below.

2.1 The Provincial Policy Statement, 2020, 2024

The Provincial Policy Statement (PPS) was issued under Section 3 of the Planning Act (Government of Ontario, 1990b). The current PPS came into effect May 1, 2020 (Government of Ontario, 2020). Natural features are afforded protections under Section 2.1 of the PPS. Protections may include maintenance, restoration, and improved function of diversity, connectivity, ecological function, and biodiversity of natural heritage systems. These protections restrict development and site alteration in significant natural areas (e.g., woodlands, wetlands, wildlife habitat) unless it can be demonstrated that there will be no negative effects on the features and ecological functions of those natural areas. Technical guidance for implementing the natural heritage policies of the PPS is found within the second edition of the Natural Heritage Reference Manual for Natural Heritage Policies of the Provincial Policy Statement, 2005 (NHRM; MNR, 2010). Importantly, while the 2020 PPS was the version in effect at the start of this study, it must be noted that the Province has approved the updated Provincial Planning Statement 2024 (i.e. as of






Figure 1 Site Location

Legend

 Site Boundary



Project: TAGG 1672.1
Map File: TAGG 1672 Map 2410B.map
Universal Transverse Mercator - Zone 18 (N)
Printed on: 2024-10-22



August 20, 2024); it came into effect on October 20, 2024. The revised PPS is intended to simplify and integrate existing policies to achieve housing objectives while providing tools for municipalities to deliver on housing objectives. While the 2024 edition will formally be the planning document in effect going forward, other than renumbering the relevant policies, there have been no meaningful changes related to Natural Heritage considerations between the two versions.

2.2 The City of Ottawa Official Plan (2021)

The City of Ottawa Official Plan (OP; City of Ottawa, 2021) was updated and recently approved by the Ministry of Municipal Affairs and Housing as part of a comprehensive review. Pursuant to subsections 17(36.5) and (38.1) of the Planning Act, the decision of the Minister of Municipal Affairs and Housing regarding an official plan adopted in accordance with section 26 of the Planning Act is final and not subject to appeal. Accordingly, the new City of Ottawa Official Plan, as approved with modifications by the Minister, came into effect on November 4, 2022. The OP provides a vision for the future growth of the city and a policy framework to guide the city's physical development. With respect to natural heritage considerations addressed under an EIS, the OP provides a framework through which species at risk and other wildlife (and their habitats), forested areas, wetlands and surface water features must be reviewed. Key portions of the OP to be considered include:

The Environmental Impact Study Guidelines (City of Ottawa, 2023) - which outlines study requirements of the EIS;

OP Schedule C11 - which outlines the Natural Heritage System Features overlay and Natural Heritage System Core Areas;

OP Section 4.8.1 - under which the City recognizes the following natural heritage features, as defined in Ottawa's Environmental Impact Study Guidelines:

- a) Significant wetlands;
- b) Habitat for endangered and threatened species;
- c) Significant woodlands;
- d) Significant valleylands;
- e) Significant wildlife habitat;
- f) Areas of Natural and Scientific Interest;
- g) Urban Natural Features;
- h) Natural Environment Areas;
- i) Natural linkage features and corridors;
- j) Groundwater features;
- k) Surface water features, including fish habitat;
- l) Landform features; and
- m) Natural features or natural areas having significant cultural, economic, or historical value to the Algonquin Anishinabe Host Nation.

Significant Woodlands: Guidelines for Identification, Evaluation, and Impact Assessment (City of Ottawa, 2022b) - which identifies wooded areas within the urban boundary that are >0.8 ha and have been continuously forested for > 60 years as "Significant Woodland";



OP Section 4.9.3 – which provides guidelines for development and site alteration near surface water features through the provision of minimum setbacks and directives to retain wetland areas and the requirement to complete headwater drainage feature assessments (HDFA) to provide management recommendations for headwater features; and

The Protocol for Wildlife Protection during Construction (City of Ottawa, 2022a) – which identifies best management practices to be employed through construction to reduce the direct impacts of development on wildlife.

2.3 Conservation Authorities Act, 1990

Conservation Authorities were created to address erosion, flooding, and drought concerns regionally by managing at the watershed level. Conservation Authorities were given the ability to regulate under Section 28 of the Conservation Authorities Act (Government of Ontario, 1990a). The Act obliges Conservation Authorities to implement Ontario Regulations 42/06 and 146/06 to 182/06 Regulation of Development, Interference with Wetlands and Alterations to Shorelines and Watercourses under Section 28 of the Conservation Authorities Act for relevant works. This project falls under the jurisdiction of the Rideau Valley Conservation Authority (RVCA).

Bill 23, which was passed on November 28th, 2022, and received Royal Assent the same day, introduced a series of legislative and proposed regulatory changes affecting conservation authorities. It is now in effect. Among the changes under Bill 23, the definition of “watercourse” was updated from an identifiable depression to a defined channel having a bed, and banks or sides.

2.4 Species at Risk Act, 2002

The federal Species at Risk Act (SARA; Government of Canada, 2002) is administered by Environment and Climate Change Canada (ECCC) and provides direction to protect and ensure the survival of wildlife species in Canada. The purpose of the SARA is to prevent populations of wildlife from becoming Extirpated, Endangered, or Threatened, provide recovery Endangered or Threatened species, and to manage other species to prevent them from becoming Endangered or Threatened.

All species listed on Schedule 1 of SARA are afforded protection on federal lands. Aquatic species and species of migratory birds protected by the Migratory Birds Convention Act (MBCA; Government of Canada, 1994) and listed as Endangered, Threatened, or Extirpated under Schedule 1 of SARA are protected wherever they occur in Canada, regardless of land ownership. SARA protections for other species do not normally extend to privately owned land. However, the Federal Minister of ECCC can and has imposed SARA protections on private projects where habitat is deemed “...*necessary for the survival or recovery of the species...*” in the area of concern.

2.5 Endangered Species Act, 2007

The provincial Endangered Species Act (ESA; Government of Ontario, 2007) is administered by the Ministry of Environment, Conservation, and Parks (MECP) and provides protection for species at risk (SAR) and their habitat. Species listed as Endangered, Threatened, or Extirpated and their habitats (e.g., areas essential for



breeding, rearing, feeding, hibernation, and migration) are automatically afforded legal protection under the ESA.

2.6 Fisheries Act, 1985

The federal Fisheries Act (Government of Canada, 1985) is administered by Fisheries and Oceans Canada (DFO) and provides protections to fish, fish habitat, and fisheries. Specifically, the Fisheries Act in its current version provides protection for all fish and fish habitat, and prohibits the harmful alteration, disruption or destruction of fish habitat.

2.7 Migratory Birds Convention Act, 1994

Nesting migratory birds are protected under the MBCA (Government of Canada, 1994). No work is permitted that would result in the destruction of active nests (nests with eggs or young birds) or the wounding or killing of bird species protected under the MBCA and/or associated regulations (e.g., SARA).

2.8 Fish and Wildlife Conservation Act, 1997

The provincial Fish and Wildlife Conservation Act (FWCA; Government of Ontario, 1997) governs the hunting and trapping of a variety of wildlife including mammals, birds, reptiles, amphibians, and fish in Ontario, thereby facilitating the protection of wildlife and their habitat. It also provides general protections for many species otherwise covered under the MBCA, the ESA and/or SARA.

2.9 Greater Cardinal Creek Subwatershed Management Plan

The Greater Cardinal Creek Subwatershed Management Plan (City of Ottawa & AECOM, 2014; herein "GCCSMP") is a City of Ottawa document addressing land use planning and environmental issues within the subwatershed, with a specific focus on development pressures, water quality, and slope stability. The plan includes policies for protection and potential habitat restoration opportunities within the subwatershed. The plan was prepared in accordance with the City of Ottawa Official Plan Policy 2.43 – Watershed and Subwatershed Plans. The Plan specifically addresses Cardinal Creek Village (Section 2.6).

The GCCSMP provides for general management recommendations to for protection and improvement of Natural System Heritage Features (i.e. the Significant Woodland) where possible including:

- Preventing any further loss or intrusion into component features;
- Preventing any further fragmentation of linkages;
- Prevent, and/or minimize, road crossings through linkages, particularly where there are watercourses;
- Preserving smaller isolated woodlots where possible; and
- Where natural features abut rear yards, installing appropriate fencing to prevent incremental intrusion. Retaining mature trees or tree clusters.

The GCCSMP provides recommended minimum watercourse setbacks for Cardinal Creek (and its tributary which passes through the Site) as the greater of:



- a) Regulatory flood line
- b) Geotechnical limit of hazard lands
- c) 30 m from normal high-water mark
- d) 25 m from top of bank
- e) Setback as determined through an Environmental Impact Statement
- f) Setback as determined through a Drain Engineer's report.

Regulatory flood lines for this area are determined and set by the RVCA. Geotechnical limits of hazard will be determined through a geotechnical study of the site. The normal high-water mark is the mark made by the action of water under natural conditions on the shore or bank of a watercourse or waterbody, the action having been so common or usual or so long continued that it has created a distinction in the general terrestrial vegetation, in changes in soil characteristics or by the edge of some embankment particularly scored by the action of water. It is a variable line in characteristic indicators and distinctiveness, and it is identified by the consideration of all visible evidence, not alone by one indicator, as located by an Ontario Land Surveyor.

The term top of bank can refer either to top of a channel bank as in the maximum point to which water can rise within the channel before spilling across the adjacent land, or it can refer to the top of slope of the associated valley, the point up the valley side where the pitch first levels out. This latter definition is (potentially) different again from the stable top of slope referred to in the current City OP. Appendix E of the GCCSMP specifically indicates that the above list of criteria were determined from HDFA setback guidelines including:

...the greater of 30 m from the centreline, 30 m from the normal high water mark, 25 m from the top of bank, or 15 m from the top of slope...

As the identified list of criteria formally selected for inclusion in the GCCSMP is clearly identified as a subset of criteria prescribed to both "top of bank and, independently, to "top of slope", the term top of bank within the context of management plan must be interpreted as referring to top of the channel bank.

Setback requirements for headwater features that would be retained are the same. For smaller headwater channels providing indirect fish habitat, setback requirements from the top of the bank (or the centerline for the smallest headwater features) are reduced to 15 m where they are retained.

The GCCSMP requires the protection of significant valleylands, though does not specifically identify the presence of this class of natural heritage feature on the site. Other than the reference to a 15 m setback to top of slope from the HDFA guidelines, the GCCSMP does not directly provide a specific setback distance for the protection of valleys.

3.0 PROPERTY IDENTIFICATION

The Site (Figure 1) currently includes two major parcels (1296 and 1400 Old Montreal Road; 45.499° N, 75.458°W), encompassing the northern portion of each property, from a tributary of Cardinal Creek northward, comprising 56.8 ha. The Site is predominantly agricultural, with scattered tree stands and a forested valley corridor along the tributary on the south edge of the Site. The majority of the Site is zoned



Rural (RU), while a portion of the west side of the Site is zoned Rural Institutional (RI) and is currently under development.

The Site is bordered by:

- Old Montreal Road, portions of the Cardinal Creek Village development, forested lands and the Ottawa River to the north;
- Agricultural lands, rural residential properties and Cardinal Creek to the west;
- Cox County Road, rural residential properties, and forested lands to the east; and
- Agricultural lands to the south.

4.0 METHODOLOGY

4.1 Desktop and Background Data Review

4.1.1 General Records Review

Background information was obtained from online databases and geographic information system mapping applications to review relevant information. Aerial imagery from Google Earth, the SNC Geoportal and the City's geoOttawa systems was used to identify existing features and confirm information found in the background review.

4.1.2 Species at Risk Screening

The review of existing information included a preliminary SAR screening for species listed under the federal SARA and provincial ESA having some record of occurrence within the broader vicinity of the Site. The screening was completed following the *Draft Client's Guide to Preliminary Screening for Species at Risk*. The results of the screening process informed the list of species that were considered in the assessment of the potential for development impact(s) to SAR or SAR habitat.

Where it is determined through the EIS process that there is an anticipated impact of the development on SAR, an Information Gathering Form (IGF) is typically submitted to MECP for further review. The IGF process, however, is not generally necessary where the SAR management process may be handled through a Notice of Activity process associated with the Ontario Conservation Fund under O.Reg. 829/21.

On-line databases queried for SAR, provincially rare species, and natural heritage features included the following:

- Species at Risk in Ontario (SARO; Ministry of Environment, Conservation, and Parks (MECP, 2024);
- Species at Risk Public Registry (Government of Canada, 2024);
- Natural Heritage Information Centre (NHIC; Ministry of Natural Resources, and Forestry (MNRF, 2024c);



- Land Information Ontario (MNRF, 2024b);
- Aquatic Species at Risk Map (DFO, 2023);
- Ontario Reptile and Amphibian Atlas (Ontario Nature, 2019);
- Ontario Breeding Birds Atlas (Birds Canada et al., 2009);
- Ontario Butterfly Atlas (Toronto Entomologists' Association, 2024);
- eBird (The Cornell Lab of Ornithology, 2024);
- iNaturalist (California Academy of Sciences and National Geographic Society, 2024);
- Bumble Bee Watch (Wildlife Preservation Canada et al., 2024);
- Recovery Strategy for the Little Brown Myotis (*Myotis lucifugus*), Northern Myotis (*Myotis septentrionalis*), and Tri-colored Bat (*Perimyotis subflavus*) in Ontario (Humphrey & Fotherby, 2019);
- Recovery Strategy for the Eastern Small-footed Myotis (*Myotis leibii*) in Ontario (Humphrey, 2017); and
- Fish ON-Line (MNRF, 2024a).

4.1.3 Site-Specific Background Data Review

In addition to the general background data outlined above, site-specific background data were obtained through previously prepared reports. The following reports were reviewed:

- Fluvial Geomorphological and Erosion Threshold Assessment, Tributary of Cardinal Creek: 1296 and 1400 Old Montreal Road (GEO Morphix, 2024);
- Environmental Impact Statement and Tree Conservation Report: Cardinal Creek Village South Portion (Muncaster Environmental Planning Inc., 2021);
- Cardinal Creek Village – South Side: Headwater Drainage Feature Assessment (Bowfin Environmental Consulting Inc., 2021);
- Geotechnical Investigation: Proposed Residential Development – Cardinal Creek Village South (Paterson Group, 2023);
- Preliminary Geotechnical Review – Proposed SWMP (Paterson Group, 2023);
- Geotechnical Response to City Comments (Paterson Group, 2023);
- Slope Stability Assessment of Existing Slope Failure (Paterson Group, 2023); and



- Geotechnical Response to Third-Party Landslide Risk Assessment Report Review Letter (Paterson Group, 2023).

4.1.4 Agency Consultation

The Site is located within the jurisdictions of the City of Ottawa and RVCA.

4.2 Field Surveys

4.2.1 Site Work Summary

KAL Biologists completed an extensive suite of field studies through the spring and summer of 2024. Table 1 provides a summary of all field visits. Specific details of each program are further described under each study type (e.g. breeding bird surveys) in the relevant sub-sections following through the remainder of Section 4.2. Specific survey stations are shown in Figure 2.

Table 1 Field Study Dates





Date	Purpose	Conditions	Personnel
March 28, 2024	<ul style="list-style-type: none"> • HDFA #1 	<ul style="list-style-type: none"> • +8°C • Overcast • Wind 15 km/h SW 	<ul style="list-style-type: none"> • Jenni Velichka • Kesia Miyashita
May 28, 2024	<ul style="list-style-type: none"> • HDFA #2 • Electrofishing 	<ul style="list-style-type: none"> • 19°C • Partly cloudy • Wind 25 km/h W 	<ul style="list-style-type: none"> • Jenni Velichka • Kurtis Westbury
June 14, 2024	<ul style="list-style-type: none"> • Breeding bird survey #1 	<ul style="list-style-type: none"> • 17°C • Sunny • Wind 10 km/h NW 	<ul style="list-style-type: none"> • Maren Nielsen
June 20, 2024	<ul style="list-style-type: none"> • Breeding bird survey #2 • Acoustic bat monitor deployment 	<ul style="list-style-type: none"> • 28°C • Sunny • Wind 14 km/h W 	<ul style="list-style-type: none"> • Maren Nielsen
July 4, 2024	<ul style="list-style-type: none"> • Breeding bird survey #3 • Acoustic bat monitor pickup • HDFA #3 • Ecological Land Classification • Black Ash Assessment 	<ul style="list-style-type: none"> • 27°C • Sunny • Wind 10 km/h SW 	<ul style="list-style-type: none"> • Robert Hallett • Nicholas Schulz






Figure 2 Field Study Locations

Legend

-  Site Boundary
-  Acoustic Bat Monitor
-  Breeding Bird Survey Station
-  HDFs Electrofished

N



Project: TAGG 1672.1
 Map File: TAGG 1672 Map 2410B.map
 Universal Transverse Mercator - Zone 18 (N)
 Printed on: 2024-10-22



4.2.2 Surface Water Characterization

Aerial imagery and public databases were reviewed to determine wetland areas and watercourses (City of Ottawa, 2024; MNRF, 2024c; Rideau Valley Conservation Authority, 2023). Any wetlands on the Site were delineated and characterized in the field as part of the Ecological Land Classification (ELC) exercise (see Section 4.2.3 below). A Headwater Drainage Feature Assessment (HDFA) was conducted for the Site following the methods per the *Evaluation, Classification and Management of Headwater Drainage Features Guidelines* (Toronto and Region Conservation Authority & Credit Valley Conservation, 2013).

The HDFA protocol requires up to three surveys of HDFs on a site. The first is conducted near the spring freshet to identify channel and wetted dimensions at peak water levels. Fish communities and habitats are assessed later in the spring for those HDFs hydrologically capable of supporting fish. Water levels of features not found to be dry during the second visit are checked once more in mid to late summer to assess their status as permanent watercourses. HDFA surveys were completed on March 28, May 28, and July 4, 2024.

The Standard level of assessment follows Ontario Stream Assessment Protocol (OSAP) methodologies for descriptions of flow conditions, riparian vegetation and site features that are important components of habitat (headwater sampling protocol OSAP S4.M10) and includes an electrofishing survey to describe fish and fish habitat (OSAP S4.M10). Additionally, the Ecological Land Classification for Southern Ontario (ELC) was applied to the Site (Lee et al., 1998), with specific focus on the riparian zone of each segment and determined habitat community types present on the Site.

4.2.3 Ecological Land Classification

Vegetation communities on the Site were identified and mapped in the field on July 4, 2024, using standard Ecological Land Classification (ELC) methods for Ontario (Lee et al., 1998). This method provides a consistent approach to identify, describe, and map vegetation communities or physiographic features on the landscape based on dominant plant species and soil composition. This method results in a standardized description of each vegetation community to capture the natural diversity and variability of communities within a site and to provide insight into available habitat and the type of species that may be present. More specifically, the classifications from ELC provide a basis for determining whether potential habitat for a given SAR or other ecological value may be present.

A desktop review of available aerial imagery and preliminary field visits informed how the Site generally divides into vegetation communities based on variation in land cover, topography, and vegetation structure. The dominant plant species were recorded within each proposed ecosite in the field to further divide ecosites into vegetation types (the finest resolution in ELC), where possible. Soil samples were taken using a 120 centimeter (cm) long soil auger to characterize community substrates. Representative photos of each ELC unit on the Site were taken and are included with the community descriptions in this report.

4.2.4 Tree Inventory, Butternut and Black Ash

A detailed tree survey was performed for the Site on July 4, 2024, following the City of Ottawa TCR guidelines. Forest groupings and notable trees to be retained on and adjacent to the Site were documented, characterized and mapped. Butternut (*Juglans cinerea*) and Black Ash (*Fraxinus nigra*) trees (both Endangered under the ESA) were specifically searched for. Any individuals of those species encountered were fully



characterized to meet provincial requirements, and formal Black Ash Assessments were undertaken for any Black Ash trees onsite.

4.2.5 Breeding Birds

Morning breeding bird surveys were performed using point counts following the Ontario Breeding Bird Atlas Guide for Participants (Ontario Breeding Bird Atlas, 2001). Breeding bird surveys are to be completed from survey stations that, combined, provide suitable viewing of all habitats on a site on calm weather days with light wind (less than 3 on the Beaufort Scale) and no precipitation. As per the Ontario Breeding Bird Atlas, three rounds of surveys must take place between sunrise and five hours after sunrise between May 24 and July 10. Surveys took place during the mornings of June 14, June 20, and July 4, 2024.

A total of four (4) breeding bird survey stations were established in representative habitats on the Site (Figure 2). All incidental observations were recorded while moving between survey points as well as during other visits to the Site. Birds were identified by song and/or direct visual observation.

Bird species were classed as regionally rare based on an analysis of data from the Atlas of Breeding Birds of Ontario (2009) based on Hill's Site Regions, now Ecoregions. The federal and provincial significance of bird species were classed based on species' listings under Schedule 1 of SARA and the ESA, and species tracked by NHIC (MNRF, 2023c; for non-SAR species considered provincially significant).

4.2.6 Bats and Other Mammals

Bat monitoring was completed following acoustic surveys under the MNRF's Survey Protocol for Species at Risk Bats within Treed Habitats (2017). This is currently the recommended protocol for confirming the presence/absence of Little Brown Myotis (*Myotis lucifugus*), Northern Myotis (*Myotis septentrionalis*), and Tri-coloured Bat (*Perimyotis subflavus*), where it is determined that potentially suitable habitat for the establishment of maternity roosts is present. Wooded areas on the Site were deemed potentially suitable habitat for the establishment of maternity roosts during KAL's preliminary desktop review and initial field visits. Trees with characteristics suitable for bat roosting were observed in the area.

All species of bats in a given area are detectable under this protocol if ultrasonic acoustic monitors are used and the signal-to-noise ratio can be analyzed from sonogram displays to identify bat calls to species level. Under the protocol, acoustic monitors are to be installed for a minimum of 10 nights between June 1 and June 30, with recordings commencing after dusk and continuing for five hours. KAL installed one acoustic monitor on the Site (Figure 2). The acoustic monitor was placed in this location to capture the best potential bat habitat on the Site (potential roosting habitat in wooded areas and potential foraging habitat over adjacent open areas) and to increase the likelihood of detecting bats based on their echolocating behaviour. Bats use echolocation more frequently in cluttered environments (Falk et al., 2014), so installing monitors along the edges of wooded areas rather than in the middle of open foraging areas likely increases bat detectability. The monitor was installed on June 20, and removed on July 4, 2024 (14 nights of data collection).

Incidental observations of other mammals present in the Study Area were collected during all field visits. Mammal observations were limited to sightings of scat, tracks, and in some cases, direct observations.



5.0 RESULTS

Site existing conditions are mapped and shown in Figure 3.

5.1 General Natural Heritage Context

The nearest lands zoned EP-Environmental Protection surrounding the Site are approximately 750 m west of the west edge of the Site (the Cardinal Creek valley) and approximately 650 m north of the north edge of the Site (forest adjacent to Cardinal Creek Village Phase 7). The closest provincially significant wetland is the Petrie Island Wetland, located approximately 1.9 km to the northwest of the Site. The nearest Life Science Area of Natural and Scientific Interest (ANSI) is also associated with Petrie Island. An Earth Science Area of Natural and Scientific Interest is located along Cardinal Creek, comprising limestone karst formations conveying the creek flow underground for approximately 250 m. This feature is approximately 1.1 km south of the southwest corner of the Site.

Significant valleylands and significant woodlands are associated with the south tributary, along the south edge of the Site. Significant natural heritage features are discussed further in Section 5.7 below.

5.2 Landforms, Soils, and Geology

The Site topography is characterized as gently sloping from east to west. Soils were characterized as topsoil over stiff silty clay and glacial till. Areas of fill were documented on the north edge of the Site along Old Montreal Road. Relatively steep slopes occurred along the tributary to Cardinal Creek at the south edge of the Site; the slopes ranged from 3 to 15 m high and were generally 5H:1V, with localized sections of 1H:1V. The slopes comprised stiff, brown silty clay. Some toe erosion was observed along the valley wall adjacent to the tributary.

Regionally, soils in the vicinity of the Site are characterized as a mosaic comprising the Rideau, Grenville and Farmington soil associations. Soils were characterized as gently sloping to nearly level with good to imperfect drainage. Portions of Cardinal Creek and the south tributary on-Site were characterized as Eroded Channels, with narrow creek beds and steep valley walls with slopes greater than 15%.

5.3 Surface Water, Groundwater and Fish Habitat

5.3.1 General Context

The Site is located within the Cardinal Creek subwatershed (City of Ottawa & AECOM, 2014). Cardinal Creek is located approximately 600 m from the west of the Site; a tributary to Cardinal Creek is located along the south edge of the Site. The tributary traverses the Site from east to west.

Farm ditches (Reaches 1-4 and 6-7) function within an agricultural context that would change substantially with urban development. These features were not noted within the GCCSMP and would be either removed or subject to significant alteration/realignment as part of a community design. Setback requirements are thus not identified in this ECR for small farm ditches.



















Figure 3 Existing Conditions

Legend

ELC Ecosites

-  CUM1-1
-  CV
-  FODM5-9
-  FODM6-5
-  MEGM3-8
-  OAGM1
-  THDM4-1
-  WODM4

-  HDF/
Watercourse
-  Top of Bank
-  Top of Slope
-  Fish Barrier
-  Black Ash



0 200 m

Project: TAGG 1672.1
 Map File: TAGG 1672 Map 2410B.map
 Universal Transverse Mercator - Zone 18 (N)
 Printed on: 2024-10-22



For Reach 5 (identified for “Protection” per the GCCSWS) and its downstream continuation as the Cardinal Creek Tributary, setback considerations as indicated in the GCCSWS (per Section 2.9 above) are the same and consider both the normal high water mark and top of bank. For these reaches, however, the normal high water mark and top of bank (i.e. of the channel) are effectively equivalent. The setback to the watercourse is thus the greater of the geotechnical limit of hazard lands or 30 m from the normal high water mark. The geotechnical limit of hazard lands will be determined separately through appropriate geotechnical studies. Setbacks from the Cardinal Creek Tributary valley are discussed separately below (Section 5.7.2).

Per the GCCSMP, no sensitive groundwater recharge areas extend onto the Site. “Watercourse 76” as indicated in Figure 2.3 of the GCCSMP was not observed to exist as a permanent watercourse crossing the entire central portion of the Site from east to west. While parts of Reaches 1, 4 and 7 align disjointedly with portions of Watercourse 76 as mapped, the mapping of Watercourse 76 as a contiguous permanent stream in the GCCSMP must be considered outdated and no longer relevant to future Site development.

5.3.2 Headwater Drainage Features

A Headwater Drainage Feature Assessment (HDFA) was completed in 2021 (Bowfin Environmental Consulting Inc., 2021). KAL completed a field review and update in spring and summer of 2024. The HDFA identified eight (8) HDFs located on the Site, the majority of which were characterized as farm drains (Figure 3). The results of the HDFA are described below.

5.3.2.1 General Reach Descriptions

Reach 1

Reach 1 is a channelized/constrained swale feature located in the eastern portion of the Site, flowing from east to west originating at Cox Country Road. Reach 1 was observed to have minimal flow during the spring freshet, with narrow-leaved emergent and shrub in-stream vegetation and a silty organic substrate.

Reach 2

Reach 2 is a channelized/constrained feature and farm drain flowing southward from Reach 1. This reach was observed to have minimal flow during spring freshet, and contain shrub and narrow-leaved emergent in-stream vegetation, with a silty clay substrate.

Reach 3

Reach 3 is a channelized/constrained feature that flows westward from Reach 2, following the boundary of the WODM4 vegetation community, and turning southward flowing into the Cardinal Creek tributary (Reach 5). Reach 3 had minimal flow during spring freshet, contained shrub and narrow-leaved emergent in-stream vegetation, and has a clay organic substrate.

Reach 4

Reach 4 originates from the upper portion of Reach 2, and traverses westward, turning southward and flowing into the Cardinal Creek tributary (Reach 5). Reach 4 was observed to have minimal flow during spring



freshet, with predominantly narrow-leaved emergent in-stream vegetation, with some shrubs present. Reach 4 has a silty clay substrate.

Reach 5

Reach 5 constitutes the upper reach of the tributary to Cardinal Creek. Combined, they traverse the Site from east to west, originating near Cox Country Road and Jonquille Way. As the watercourse descends into the valley, it is considered a permanent stream per the GCCSWS. Reach 5 is a defined natural channel with substantial baseflow observed. Robust emergent, shrub, narrow and broad-leaved emergent, trees, and herbaceous in-stream vegetation were observed throughout the reach. Clay, silt, and cobble substrates were observed.

Reach 6

Reach 6 is a swale feature that flows southward, that was found to be dry during spring freshet. Tree and shrub vegetation were observed within this reach. A few isolated wetland pockets of standing water were observed, with silty organic substrate.

Reach 7

Reach 7 is a swale feature that flows east to west originating in the open agricultural area (OAGM1) in the northern portion of the Site, outletting to the meadow in the northwest site corner. Reach 7 was observed to have standing water during spring freshet, and have robust and narrow-leaved emergent vegetation, shrubs, and trees in-stream. Organic clay substrates are present within this reach.

Reach 8

Reach 8 is a roadside ditch feature along Old Montreal Road in the northwestern site corner, east of the construction access gate. It was observed to have standing water during the spring freshet, with silty organic substrate and broad-leaved emergent, robust emergent, narrow-leaved emergent, shrub and tree in-stream vegetation.

5.3.2.2 Component Classifications

Tables 2-5 below summarize the functions provided by the eight (8) drainage features.



Table 2 Hydrology Classifications for HDFs

Drainage Feature	Hydrology Classification					
	Assessment Period	Flow Conditions		Flow Classification	Modifiers	Hydrological Function
		Description	(OSA P Code)			
1	March 28, 2024	Minimal Surface flow	4	Ephemeral	No source other than spring run-off and after heavy rain	Contributing Functions
	May 28, 2024	Minimal Surface flow	4			
2	March 28, 2024	Minimal Surface flow	4	Ephemeral/Intermittent	No source other than spring run-off and after heavy rain	Contributing Functions
	May 28, 2024	Minimal Surface flow	4			
3	March 28, 2024	Minimal Surface Flow	4	Ephemeral	No source other than spring run-off and after heavy rain	Contributing Functions
	May 28, 2024	Minimal Surface flow	4			
4	March 28, 2024	Minimal Surface flow	4	Intermittent	No source other than spring run-off and after heavy rain	Valued Functions
	May 28, 2024	Minimal Surface flow	4			
5	March 28, 2024	Substantial Flow	5	Perennial	Water is present throughout the year. Upstream minimal flow significant flow downstream in valley	Important Functions
	May 28, 2024	Substantial Flow	5			
6	March 28, 2024	Minimal Surface flow	4	Ephemeral	No source other than spring run-off and after heavy rain	Contributing Functions
	May 28, 2024	Dry	1			
7	March 28, 2024	Dry	1	Dry or Standing Water	No source other than spring run-off and after heavy rain	Limited Functions
	May 28, 2024	Dry	1			
8	March 28, 2024	Standing Water	2	Ephemeral	No source other than spring run-off and after heavy rain	Contributing Functions
	May 28, 2024	Dry	1			



Table 3 Riparian Classifications

Drainage Feature	Riparian Classification			
	OSAP Descriptions	OSAP Riparian Codes	ELC Codes	Riparian Conditions
1	RUB - Scrubland	RUB - 5	THDM4-1, OAGM1	Important/Valued Functions
	LUB - Meadow	LUB - 4		
2	RUB - Cropped	RUB - 3	OAGM1	Limited Functions
	LUB - Cropped	LUB - 3		
3	RUB - Forest	RUB - 7	WODM4, OAGM1	Important Functions
	LUB - Cropped	LUB - 3		
4	RUB - Cropped	RUB - 3	OAGM1	Limited Functions
	LUB - Cropped	LUB - 3		
5	RUB - Forest	RUB - 7	FODM6-5	Important Functions
	LUB - Forest	LUB - 7		
6	RUB - None	RUB - 1	CV	Limited Functions
	LUB - Meadow	LUB - 4		
7	RUB - Cropped	RUB - 3	OAGM1, CUT1-1	Limited Functions
	LUB - Cropped	LUB - 3		
8	RUB - Meadow	RUB - 4	CUT1-1	Valued Functions
	LUB - Meadow	LUB - 4		

Table 4 Fish and Fish Habitat Classification

Drainage Feature	Fish Habitat Classification		
	Fish Observation	Fish & Fish Habitat Designation*	Modifiers/Notes
	Fishing effort		
1	No fish present; 195 S	Contributing Functions	
2	Dry	Contributing Functions	
3	Dry	Contributing Functions	
4	No fish present; 710.8 S	Contributing Functions	
5	No fish present (upper reach); 526.6 S	Valued Functions	Upper reach electrofished; no fish present. Lower reach in the valley was not electrofished as fish habitat is assumed present
6	Dry	Contributing Functions	
7	Dry	Contributing Functions	
8	Dry	Contributing Functions	



Table 5 Terrestrial Classifications

Drainage Feature	Terrestrial Classification		
	Description	Amphibians	Terrestrial Classification
1	This reach is a swale that is heavily vegetated and holds water for much of the summer. There is no wetland habitat present.	No frogs were observed within the vicinity of this feature	Limited Functions
2	This reach is a channelized feature that conveys flow into Reach 5. There is no wetland habitat present.	No frogs were observed within the vicinity of this feature	Limited Functions
3	This reach is a channelized feature that conveys flow into Reach 5. There is no wetland habitat present.	No frogs were observed within the vicinity of this feature	Limited Functions
4	This reach is a channelized farm drain holds water for much of the summer and conveys flow to Reach 5.	No frogs were observed within the vicinity of this feature	Limited Functions
5	This is a permanent stream feature and tributary to Cardinal Creek. Forest provides important riparian habitat connecting the Site with downstream features. No wetland habitat is present.	No frogs were observed within the vicinity of this feature	Contributing Functions
6	This reach is a swale that is dry for most of the year. There is no wetland habitat present.	No frogs were observed within the vicinity of this feature	Limited Functions
7	This reach is a swale that is dry for most of the year. There is no wetland habitat present.	No frogs were observed within the vicinity of this feature	Limited Functions
8	This reach is a roadside ditch that is dry for most of the year. There is no wetland habitat present.	No frogs were observed within the vicinity of this feature	Limited Functions

5.4 Ecological Land Classification

A total of eight (8) distinct landcovers or ELC units were delineated on the Site (Figure 3). The majority of the Site is dominated by an Open Agriculture – Annual Row Crops (OAG) ecosite. The valleylands associated with the Cardinal Creek tributary are characterized by a Fresh – Moist Sugar Maple – Hardwood Deciduous Forest Type (FODM6-5), a Dry - Fresh Deciduous Woodland Ecosite (WODM4), and a Reed Canary Grass Graminoid Meadow Type (MEGM3-8). Three isolated vegetated areas within the OAG ecosite are characterized by a Native Deciduous Regeneration Thicket Type (THDM4-1) adjacent to the eastern Site boundary and within the northeastern corner adjacent to Old Montreal Road and Cox Country Road, and a Dry – Fresh Sugar Maple – Hardwood Deciduous Forest Type (FODM5-9) ecosite in an isolated forest stand in the northern portion of the Site. The western portion of the Site is characterized by a Cultural Thicket (CUT1-1) ecosite and a large area of fill piles and construction material (Constructed - CV).



5.4.1 Open Agriculture – Annual Row Crops (OAGM1)

The Open Agriculture – Annual Row Crops is planted with corn crops. Based on a review of historic aerial imagery, this area has been actively farmed since at least 1976 (City of Ottawa, 2024).

5.4.2 Fresh – Moist Sugar Maple – Hardwood Deciduous Forest Type (FODM6-5)

The valleylands, riparian area, and upland areas associated with the Cardinal Creek tributary is dominated by a Fresh – Moist Sugar Maple – Hardwood Deciduous Forest Type (FODM6-5). The tree canopy is dominated by Sugar Maple (*Acer saccharum*), with American Beech (*Fagus grandifolia*), Trembling Aspen (*Populus tremuloides*), White Birch (*Betula papyrifera*) and Eastern Hemlock (*Tsuga canadensis*) within the shady valley bottom adjacent to the tributary. No Black Ash (*Fraxinus nigra*) or Butternut (*Juglans cinerea*) trees were observed within this ecosite (Figure 3). Little to no understory and groundcover vegetation is present within this ecosite. The dominant tree species size ranged from 30-40 DBH, and 11-20 m in height, on average.

Soils within the FODM6-5 community were found to be mineral material, with a loamy A horizon extending from 0-20 cm in depth, with a sandy loam B horizon extending to a depth of 75 cm, and clay soils with a blocky structure within the C horizon (75+ cm). The exposed soil profile is shown in Figure 5 below.





Figure 4 Fresh – Moist Sugar Maple – Hardwood Deciduous Forest Type (FODM6-5)





Figure 5 FODM6-5 Soil Core

5.4.3 Dry – Fresh Deciduous Woodland Ecosite (WODM4)

East of the tributary valleylands, adjacent to the tributary and MEGM3-8 community (described below), transitioning to the OAGM1 area is characterized by a Dry – Fresh Woodland Ecosite (WODM4) vegetation community. The upper canopy of the woodland is dominated by Bur Oak (*Quercus macrocarpa*) and Red Maple (*Acer rubrum*). The subcanopy is dominated by Green Ash (*Fraxinus pennsylvanica*), Eastern Black Walnut (*Juglans nigra*), Basswood (*Tilia americana*), and Hawthorn (*Crataegus spp.*). Groundcover species include Red Raspberry (*Rubus idaeus*), Jewelweed (*Impatiens capensis*), and Poison Ivy (*Rhus radicans*). Many dead fallen trees and open groundcover (bare soil) was observed within this community. Dominant tree species ranged from 35-50 DBH in size, and 11-20 m in height, on average. Subdominant tree species ranged from <10-15 DBH on average. Three Black Ash trees were observed within this community, and are addressed further in Section 5.6.2 of this ECR.

Soils within this community were found to be moist mineral clay loam soils. Undecomposed organic material was present within the A horizon (0-5 cm), and light mottling was encountered around a depth of 30 cm within the B horizon (5 – 60 cm). A heavy clay layer was encountered around a depth of 80 cm within the C horizon (60+ cm). Soil core within the WODM4 in Figure 7.





Figure 6 Dry – Fresh Woodland Ecosite (WODM4)





Figure 7 Soil Core in WODM4

5.4.4 Reed Canary Grass Graminoid Meadow Type (MEGM3-8)

A Reed Canary Grass Graminoid Meadow Type (MEGM3-8) is associated with the upper reach of the tributary to Cardinal Creek. It is bordered by the WODM4 vegetation community to the north and south. The MEGM3-8 community is dominated entirely by Reed Canary Grass (*Phalaris arundinacea*). No trees, shrubs, or other groundcover vegetation species are present within the community.

Soils within this community were found to be mineral material, with a clay loam texture throughout all exposed horizons. Light mottling and gleying was encountered around a depth of 40 cm (Figure 9).





Figure 8 Reed Canary Grass Graminoid Meadow Type (MEGM3-8)





Figure 9 MEGM3-8 Soil Core

5.4.5 Native Deciduous Regeneration Thicket Type (THDM4-1)

A Native Deciduous Regeneration Thicket Type (THDM4-1) community is located within the eastern and northeastern portions of the Site, associated with HDF 1 (Figure 3), and at the corner of Old Montreal Road and Cox Country Road. This community is dominated by Manitoba Maple with Trembling Aspen (*Populus tremuloides*), Staghorn Sumac (*Rhus typhina*), Glossy Buckthorn (*Rhamnus frangula*), and American Plum (*Prunus spp.*). Large dead Ash trees were observed.





Figure 10 Native Deciduous Regeneration Thicket Type (THDM4-1)

5.4.6 Dry – Fresh Sugar Maple – Hardwood Deciduous Forest Type (FODM5-9)

A Dry – Fresh Sugar Maple – Hardwood Deciduous Forest Type (FODM5-9) characterizes an isolated forest stand within the northern portion of the Site, southwest of the existing home along Old Montreal Road. The upper canopy is dominated by American Elm (), Sugar Maple, Green Ash, and Basswood trees. Canopy trees ranged from 30-40 DBH on average and 11-20 m in height. The subcanopy of the FODM5-9 community was composed primarily of saplings of the same canopy species.



Soils within this community were composed mineral material, with loamy A horizon extending from 0 cm to a depth of 15 cm, and a clay loam B horizon from 15-40 cm, and a heavy silty clay C horizon (40+ cm). The exposed soil profile is shown in Figure 12.




Figure 11 Dry – Fresh Sugar Maple – Hardwood Deciduous Forest Type (FODM5-9)





Figure 12 FODM5-9 Soil Core

5.4.7 Cultural Meadow Ecosite (CUM1-1)

A Cultural Meadow Ecosite (CUM1-1) is located on the western side of the Site, south of Old Montreal Road. This area consisted of active agricultural fields until 2017. Beginning in 2018, most of the ecosite was subject to various ground works with fill piles moved around the area. The ecosite is currently dominated by Kentucky Bluegrass (*Poa pratensis*) with Canada Goldenrod (*Solidago canadensis*), Common Dandelion (*Taraxacum officinale*), and Virginia Creeper (*Parthenocissus quinquefolia*). Scattered young Manitoba Maple and Staghorn Sumac saplings have begun sprouting randomly across the area.





Figure 13 Cultural Meadow (CUM1-1) adjacent to Open Agriculture (OAGM1-1)

5.4.8 Constructed (CV)

A large area of gravel fill, construction materials and fill piles is located in the western portion of the Site (Figure 14).





Figure 14 Constructed (CV)

5.5 Wildlife Surveys

5.5.1 Breeding Birds

Breeding bird surveys were completed via three rounds of surveys on June 14, June 20, and July 4, 2024. Species observed, the station observed, and breeding evidence is shown in Table 2 below. 27 bird species were observed, belonging to common, widespread species. One at-risk bird was observed, the Eastern Wood-Pewee (Special Concern).



Table 6 Breeding Bird Survey Data

Species Observed	Station	Highest Breeding Evidence	Species Observed	Station	Highest Breeding Evidence
American Crow (<i>Corvus brachyrhynchos</i>)	BBS1, BBS 2, BBS3	Possible	Indigo Bunting (<i>Passerina cyanea</i>)	BBS1	Probable
American Goldfinch (<i>Spinus tristis</i>)	BBS1, BBS 2, BBS3	Observed	Killdeer (<i>Charadrius vociferus</i>)	BBS1	Possible
American Redstart (<i>Setophaga ruticilla</i>)	BBS3	Possible	Mallard (<i>Anas platyrhynchos</i>)	BBS2	Transient
American Robin (<i>Turdus migratorius</i>)	BBS1, BBS 2, BBS3	Probable	Northern Cardinal (<i>Cardinalis cardinalis</i>)	BBS2	Possible
Black-and-white Warbler (<i>Mniotilta varia</i>)	BBS1	Possible	Northern Flicker (<i>Colaptes auratus</i>)	BBS2, BBS3	Probable
Black-capped Chickadee (<i>Poecile atricapillus</i>)	BBS1, BBS2	Probable	Red-eyed Vireo (<i>Vireo olivaceus</i>)	BBS1, BBS3	Possible
Blue Jay (<i>Cyanocitta cristata</i>)	BBS1	Probable	Red-tailed Hawk (<i>Buteo jamaicensis</i>)	BBS2	Possible
Cedar Waxwing (<i>Bombycilla cedrorum</i>)	BBS1, BBS2	Possible	Red-winged Blackbird (<i>Agelaius phoeniceus</i>)	BBS1, BBS2, BBS3, BBS4	Observed
Chestnut-sided Warbler (<i>Setophaga pensylvanica</i>)	BBS2	Possible	Song Sparrow (<i>Melospiza melodia</i>)	BBS1, BBS2, BBS3, BBS5	Probable
Chipping Sparrow (<i>Spizella passerina</i>)	BBS4	Possible	Swamp Sparrow (<i>Melospiza georgiana</i>)	BBS4	Probable
Common Grackle (<i>Quiscalus quiscula</i>)	BBS1, BBS4	Observed	Turkey Vulture (<i>Cathartes aura</i>)	BBS1, BBS3	Transient
Common Yellowthroat (<i>Geothlypis trichas</i>)	BBS1	Probable	Wild Turkey (<i>Meleagris gallopavo</i>)	BBS2, BBS3	Probable
Eastern Wood-Pewee (<i>Contopus virens</i>)*	BBS2	Probable	Yellow Warbler (<i>Setophaga petechia</i>)	BBS2, BBS3	Probable
Great Crested Flycatcher (<i>Myiarchus crinitus</i>)	BBS1, BBS2	Probable			

* Species listed as-risk in Ontario

5.5.2 Bats and Other Mammals

One acoustic bat monitor was installed for 14 nights and placed facing an open agricultural community (OAGM1), just north of the FODM6-5 community, where the greatest likelihood for bat activity would occur on the Site. Conditions were ideal with mainly clear or cloudy nights and warm temperatures ($\geq 15^{\circ}\text{C}$). Bat species identified within the Site include Big Brown Bat (*Eptesicus fuscus*), Hoary bat (*Lasiurus cinereus*), Silver-haired Bat (*Lasionycteris noctivagans*), Tri-colored Bat (*Perimyotis subflavus*), Eastern Red Bat (*Lasiurus borealis*), and Little Brown Bat (*Myotis lucifugus*). Bat monitor location is shown in Figure 2.



Table 7 Acoustic Bat Survey Data

Survey Station	Survey Dates	Habitat Description	Big Brown Bat	Eastern Red Bat	Hoary Bat	Eastern Small-footed Bat	Little Brown Bat	Silver-haired Bat	Tri-Colored Bat	Northern Long-eared Bat	Mean Number of Calls per Night
AM-1	2023-06-20 to 2023-07-04	Open agricultural field adjacent to FODM6-5 ecosite	57	2	288	0	1	362	7	0	9

5.6 Species at Risk

An assessment of species listed under SARA and ESA was completed to identify species having some potential to occur on or near the Site, including Extirpated, Endangered, Threatened, and Special Concern species. Species listed as Extirpated, Endangered, and Threatened are afforded species and habitat protection under the ESA. Federal protections under SARA are always in force for listed species of fish and migratory birds. For species of other groups, SARA normally only applies on federal lands or on projects having some level of participation with or oversight by the federal government. However, SARA-based protections can be imposed by ministerial order on a case-by-case basis in situations where provincial-level protections are deemed inadequate to otherwise protect a species. Such protections are not expected to apply to the Site.

The SAR assessment evaluated whether the Site may provide suitable habitat for SAR (i.e. considering species known to occur in the Ottawa area; Appendix B) and whether they have potential to interact with future development of the Site. An assessment of the potential for SAR and their potential habitat was completed based on the results of the field surveys, ELC (i.e., habitat availability), and a desktop review that considered known species ranges, historic observation records, and preferred habitat requirements of these species (Appendix B). A total of 16 species subject to protections as SAR under the ESA and/or SARA were initially considered to have a moderate to high potential to occur on the Site and/or interact with the project (Table 8). Of those 16 species, three were observed to occur on the Site. Those species are discussed below.



Table 8 Species at risk with moderate or high potential to interact with the project

Species Name (Taxonomic name)	Status under Endangered Species Act	Status under Species at Risk Act (Schedule 1)	Potential to Interact with Development of the Site
Birds			
Bank Swallow (<i>Riparia riparia</i>)	Threatened	Threatened	Not detected on the Site
Bobolink (<i>Dolichonyx oryzivorus</i>)	Threatened	Threatened	Not detected on the Site
Canada Warbler (<i>Cardellina canadensis</i>)	Special Concern	Threatened	Not detected on the Site
Eastern Meadowlark (<i>Sturnella magna</i>)	Threatened	Threatened	Not detected on the Site
Eastern Wood- Pewee (<i>Contopus virens</i>)	Special Concern	Special Concern	High – detected onsite
Olive-sided Flycatcher (<i>Contopus cooperi</i>)	Special Concern	Threatened	Not detected on the Site
Wood Thrush (<i>Hylocichla mustelina</i>)	Special Concern	Threatened	Not detected on the Site
Mammals			
Eastern Red Bat (<i>Lasiurus borealis</i>)	Endangered (January 2025)	No Status	Limited/transient presence only – low probability of negative interactions if tree clearing occurs outside of the active season
Eastern Small-footed Myotis (<i>Myotis leibii</i>)	Endangered	No Status	Not detected on the Site
Hoary Bat (<i>Lasiurus cinereus</i>)	Endangered (January 2025)	No Status	Detected on the Site – migratory species, low probability of negative interactions if tree clearing occurs outside of the active season
Little Brown Myotis (<i>Myotis lucifugus</i>)	Endangered	Endangered	Limited/transient presence only – low probability of negative interactions if tree clearing occurs outside of the active season
Northern Myotis / Northern Long-eared Bat (<i>Myotis septentrionalis</i>)	Endangered	Endangered	Not detected on the Site
Silver-haired Bat (<i>Lasionycteris noctivagans</i>)	Endangered (January 2025)	No Status	Detected on the Site – migratory species, low probability of negative interactions if tree clearing occurs outside of the active season
Tri-colored Bat / Eastern Pipistrelle (<i>Perimyotis subflavus</i>)	Endangered	Endangered	Limited/transient presence only – low probability of negative interactions if tree clearing occurs outside of the active season
Vascular Plants			
Black Ash (<i>Fraxinus nigra</i>)	Endangered	No Status	High – present on the Site
Butternut (<i>Juglans cinerea</i>)	Endangered	Endangered	Low - Not observed on the Site

SAR presented in Table 8 do not include listed species that are not directly protected as SAR under the ESA or SARA (e.g. listed only as Special Concern, or are protected only federally and are not birds or fish). However, individuals of these species are protected under other regulations addressing wildlife conservation generally, such as the FWCA, the MBCA, and the PPS. In addition, species listed as Special Concern under the ESA may receive habitat protection if they are observed in habitats that meet the criteria for designation as SWH for Special Concern Species (MNRF, 2015). Species of Species Concern will be discussed with SWH in Section 5.8.



5.6.1 SAR Bats

The Committee on the Status of Species at Risk in Ontario (COSSARO) has updated the provincial status for the Hoary Bat, Silver-haired Bat, and Eastern Red Bat to Endangered. These species will receive general habitat protection on or prior to January 31, 2025. Although these species are not officially listed at the time of this ECR, it is anticipated that protections will apply throughout the development application timeline, and during future community build-out. As such, these species are considered and assessed as Endangered species in this ECR and future EIS.

The Hoary Bat and Silver-haired Bat were detected in high numbers at the monitoring stations on the Site, indicating potential roosting habitat. The Eastern Red Bat, Little Brown Myotis, and Tri-colored Bat were detected at the monitoring stations on the Site and therefore likely forage and/or roost in proximity to the Site. The numbers of detections, however, were very low, suggesting only a limited transient presence over most of the Site, with little evidence of maternal roosting activity or habitat. As Endangered species, Hoary Bat, Silver-haired Bat, Eastern Red Bat, Little Brown Myotis and Tri-colored Bat receive “general habitat protection” under the ESA. However, vegetation removal on the Site would not result in a loss of maternal roosting habitat for the Hoary Bat, Little Brown Myotis and Tri-colored Bat given the protection of the Cardinal Creek tributary lands.

Regardless, individuals of listed bat species may periodically roost diurnally in trees on the site during the active season (April 1 to September 30 inclusive; MNRF, 2017), i.e., bats could briefly use any site tree or structure as a rest stop, but only opportunistically (not as a required habitat element). Potential impacts to individual at-risk bats directly would be mitigated by clearing trees, removing structures (or commencing construction works on them) outside of the roosting season. Following this tree-clearing window would also avoid potential interactions with birds and bird nests protected under the Migratory Birds Convention Act (MBCA; Government of Canada, 1994). As such, the Hoary Bat, Silver-haired Bat, Eastern Red Bat, Little Brown Myotis and Tri-colored Bat are generally considered unlikely to be impacted by future site development. Impacts and associated mitigation measures to protect SAR bats will be outlined in a future EIS for the Site.

5.6.2 Black Ash

Black Ash (*Fraxinus nigra*), endangered under the ESA and with no status under the SARA, are a medium sized shade-intolerant hardwood tree primarily found in wetland environments like swamps, floodplains and fens. Black Ash can also occur in moist upland forests (COSEWIC, 2018). Black Ash received protection under the ESA on January 24, 2024. O.Reg 6/24 and O.Reg 7/24 set out individual and habitat protection. Black Ash habitat is defined as a radial distance of 30 m from the stem of every Black Ash that are over 8 cm at 1.37 m.

A total of three Black Ash >8 DBH were observed on the Site (Figure 3). Black Ash were located predominantly east of the tributary valleylands, within a Dry – Fresh Woodland Ecosite (WODM4) vegetation community. All three trees were determined to be healthy. Healthy trees have a canopy condition rating of 1, 2 or 3, and mortality is unlikely within five years based on severity of stressors. Unhealthy trees have a canopy condition rating on 3, 4 or 5, and mortality is expected within five years based on the severity of stressors. No Black Ash saplings were observed on the Site.

Development within any portion of this ecosites would lead to the impact or removal of healthy Black Ash protected under the ESA. Future site development that impacts healthy Black Ash will require the submission



of a Black Ash Assessment Report to the MECP and an Information Gathering Form (IGF) to support a Net Benefit Permit under the ESA.

5.7 Significant Natural Heritage Features

5.7.1 Significant Woodlands and Canopy Cover

The City of Ottawa’s (2022b) Significant Woodland Policy, defines Significant Woodlands within the urban boundary as any area 0.8 hectares in size or larger, supporting woodland 60 years of age and older at the time of evaluation. Significant Woodland on the Site was thus demarcated by delineating the boundaries of wooded areas on and adjacent to the property based on aerial imagery from 1976 (City of Ottawa, 2024). Portions of the demarcated areas that were noted as subsequently deforested in historical aerial imagery between 1976 and 2023 within the geoOttawa system were removed. Remaining areas greater than 0.8 ha in size were deemed to constitute Significant Woodland. A total of 6.0 ha of the wooded areas on the Site thus constitute Significant Woodland.

A portion of the Significant Woodlands on the Site are mapped as an Urban Natural Feature (UNF) on Schedule C11-C of the City’s OP (Figure 15; City of Ottawa, 2021).

Significant Woodland features on the site are characterized according to screening criteria per the City’s Significant Woodlands policy (2022; Table 9).

Table 9 Characterization of Significant Woodland Areas

Social Values	
Unusual recreational, educational or cultural opportunities	None. The Site consists of private property with no public use supported.
Qualifying Cultural, Heritage, or Historical Features	None. There are no existing designations within the OP.
Indigenous values established through consultation	None. No values are identified in the Greater Cardinal Creek Subwatershed Management Plan.
Hazard lands	
Constrained areas	Subject area is associated with hazards (steep or unstable slopes).
Habitat and Landscape Connectivity	
Adjacency and connectivity	Included in the Natural Heritage Features Overlay. Not part of Natural Heritage System Core Area or identified greenspace. Forested areas on the Site extend west to join Cardinal Creek and its associated forests and riparian areas.
Specialized habitat	Limited. There are no uncommon community types or rare species within the wooded areas. The current forest mix consists of trees neither especially large nor uncharacteristically old for the broader area. Three Black Ash (Endangered) were identified.











Figure 15 Significant Natural Heritage Features

Legend

-  Site Boundary
-  Significant Woodland
-  Significant Valleylands
-  Natural Heritage Features Overlay (OP Schedule C11)



Project: TAGG 1672.1
 Map File: TAGG 1672 Map 2410B.map
 Universal Transverse Mercator - Zone 18 (N)
 Printed on: 2024-10-22



5.7.1.1 iTree Canopy Assessment

An iTree Canopy assessment of the Site compares the canopy services across the Site generally. The assessment was based on distributions of 100 random sample points across the entire Site.

Table 10 Assessment of Canopy Tree Benefits on the Site

Land Cover Distribution			
Land Cover Type	General Site		
	Area (ha)	Area (%)	
Grass/Herbaceous	23.76 ± 2.42	49 ± 5.00	
Impervious Buildings	0.00 ± 0.00	0.00 ± 0.00	
Impervious Other	1.94 ± 0.97	4.00 ± 2.00	
Impervious Road	0.48 ± 0.48	1.00 ± 1.00	
Soil/Bare Ground	15.52 ± 2.26	32.00 ± 4.66	
Tree/Shrub	6.79 ± 1.68	14.00 ± 3.47	
Water	0.00 ± 0.00	0.00 ± 0.00	
Total	48.49	100%	
Tree Benefit Estimates: Carbon			
	General Site		
	Carbon (t) ± SE	CO2 Equiv. (t) ± SE	Value (CAD) ± SE
Sequestered annually in trees	20.77 ± 5.15	76.17 ± 18.88	\$5,258 ± 1,303
Total stored in trees	521.73 ± 129.31	1,912.99 ± 474.13	\$132,060 ± 32,731
Tree Benefit Estimates: Air Pollution			
Pollutant Removed Annually	General Site		
	Amount (kg) ± SE	Value (CAD) ± SE	
CO - Carbon Monoxide	6.87 ± 1.70	\$4 ± 1	
NO ₂ - Nitrogen Dioxide	34.38 ± 8.52	\$1 ± 0	
O ₃ - Ozone	365.53 ± 90.60	\$63 ± 1+	
SO ₂ – Sulfur Dioxide	34.34 ± 8.51	\$0 ± 0	
PM _{2.5} - Particulate Matter <2.5 µm	18.06 ± 4.48	\$132 ± 33	
PM ₁₀ - Particulate Matter 2.5 – 10 µm	129.89 ± 32.19	\$378 ± 94	
Tree Benefit Estimates: Hydrological			
Benefit	General Site		
	Amount (l) ±SE	Value (CAD) ± SE	
Avoided Runoff	57.02 ± 14.13	\$181 ± 45	
Evaporation	4,704.55 ± 1,166.01	N/A	
Interception	4,728.02 ± 1,171.83	N/A	
Transpiration	7,290.08 ± 1,806.83	N/A	
Potential Evaporation	35,770.41 ± 8,865.62	N/A	
Potential Evapotranspiration	35,770.41 ± 8,865.62	N/A	



5.7.2 Significant Valleylands

Significant Valleylands are defined as “valleylands with slopes greater than 15 percent and a length of more than 50 metres, with water present for some period of the year, excluding manmade features such as pits and quarries” (City of Ottawa, 2023). The Cardinal Creek tributary and associated ELC ecosites, FODM6-5 and portions of WODM4 meet the criteria to be considered significant valleylands, and constitute 3.5 ha of the Site (Figure 15). Valleylands were delineated based on field observations and available LiDAR data for the Site.

The GCCSMP does not provide a specific setback distance for the protection of Significant Valleylands. The GCCSMP, however, indicates HDFA Guidelines as the source for watercourse setback considerations, noting their identification of a general 15 m setback from the top of slope. Setbacks along Significant Valleylands should also consider geotechnical issues and wildlife habitat functions.

A 15 m setback from the existing top of slope would be situated fully outside of the Significant Woodland cover located within the Significant Valleyland feature on the Site and would sit within agricultural areas that currently do not otherwise provide wildlife habitat functions. As such, to protect the Significant Valley, the recommended setback to feature (i.e. independent of watercourse setbacks otherwise provided within the GCCSMP) is the greater of:

- a) a 15 m setback from the existing top of slope of the valley; and
- b) the geotechnical setback to the valley as determined by a geotechnical study.

5.8 Significant Wildlife Habitat

The Significant Wildlife Habitat (SWH) Criteria Schedule for Ecoregion 6E (MNRF, 2015) identifies four main types of significant wildlife habitat: seasonal concentration areas, rare vegetation communities, specialized habitat for wildlife and habitats of Species of Conservation Concern.

5.8.1 Seasonal Concentration Areas

Seasonal concentration areas include terrestrial and aquatic waterfowl stopover and staging areas, shorebird migratory stopover areas, raptor wintering areas, bat hibernacula, maternity colonies, and migratory stopover areas, turtle wintering areas, reptile hibernaculum, colonially nesting bird breeding habitat (bank and cliff; tree/shrub; ground), migratory butterfly stopover area, landbird migratory stopover areas, and deer yarding and winter congregation areas.

The background information reviewed for the Site did not identify any seasonal concentration areas for animals. No obvious signs or evidence of use as a seasonal concentration area were observed and none are likely to occur on the Site.

5.8.2 Rare Vegetation Communities or Specialized Habitat for Wildlife

Rare Vegetation Communities



Rare vegetation communities typically include those that have developed on cliff and talus slopes, sand barrens, shallow soils over limestone bedrock (alvar), old growth forests, savannahs, and tallgrass prairies. No rare vegetation communities were observed on the Site.

Specialized Wildlife Habitat

Specialized Wildlife Habitat includes waterfowl nesting areas, Bald Eagle and Osprey nesting, foraging and perching habitat, woodland raptor nesting habitat, turtle nesting areas, seeps and springs, woodland amphibian breeding habitat, wetland breeding habitat, and woodland area-sensitive bird breeding habitat.

The Red-tailed Hawk (*Buteo jamaicensis*) was observed adjacent to FODM6-5 ecosite. However, the Red-tailed Hawk is not listed as a species considered as candidate SWH for woodland raptor nesting habitat, and the Site therefore does not qualify as SWH for this category. No other specialized wildlife habitat is present on the Site.

Habitats of Species of Conservation Concern

Habitats of Species of Conservation Concern include marsh bird breeding habitat, open country bird habitat, shrub/early successional bird breeding habitat, terrestrial crayfish and special concern and rare wildlife species. Habitats of Species of Conservation Concern do not include habitats of Endangered or Threatened species as identified by the ESA. Our background review identified potential presence of three Special Concern species, including Canada Warbler, Eastern Wood-Pewee, and Olive-sided Flycatcher. Eastern Wood-Pewee was observed on the Site during breeding bird surveys. Therefore, the Site qualifies as SWH for special concern and rare wildlife species.

5.9 Other Natural Heritage Features

No Provincially Significant Wetlands (PSW) and/or Areas of Natural and Scientific Interest (ANSI) are located on or adjacent to the Site. The Site does not contain significant woodlands, valleylands or greenspace linkages. No other significant natural heritage features are located within 120 m of the Site.

6.0 OPPORTUNITIES

The Site is proposed for residential development to meet the City's approved housing projections. The design of this new residential neighbourhood provides opportunities to contribute to the urban canopy, provide natural and active recreational areas and to improve areas of aquatic habitat on site compatible with urban form and infrastructure.

Portions of the Site include retained significant forest cover associated with the Cardinal Creek tributary. Much of the Site is cleared for active agricultural purposes, and very little tree cover exists across the Site. Development on the Site has the opportunity improve canopy cover with the establishment of treed parks, open spaces, street trees, and residential trees, achieving social benefits to future residents.



HDFs on the Site currently exist in primarily open areas on the Site and are hydrologically limited, having insufficient water level through most of the year to support fish or other aquatic wildlife. The Cardinal Creek tributary (Reach 5) will be preserved and protected with a 30 m setback from the normal high water mark to protect the watercourse and significant valleylands. Future Site redevelopment is anticipated to require the construction of SWM pond facilities to support stormwater management for the area. The outlet channels for these feature(s) provide an opportunity to design local watercourses following principals of natural channel design and with increased levels of hydration that would support improved habitat for local biota beyond the limited capacity afforded by the current HDFs.



7.0 CLOSURE


This report was prepared for exclusive use by Taggart (Tamarack Homes) and may be distributed only by Taggart (Tamarack Homes). Questions relating to the data and interpretation can be addressed to the undersigned.

Respectfully submitted,

KILGOUR & ASSOCIATES LTD.



Maren Nielsen, BES, EMA
Biologist, Project Manager
E-mail: mnielsen@kilgourassociates.com
16-2285 St. Laurent Blvd, Ottawa, ON, K1G 4Z6
Office: 613-260-5555
Cell: 613-367-5562



Anthony Francis, PhD
Director of Land Development
E-mail: afrancis@kilgourassociates.com
16-2285 St. Laurent Blvd, Ottawa, ON, K1G 4Z6
Office: 613-260-5555
Cell: 613-367-5556



8.0 LITERATURE CITED

- Birds Canada, Canadian Wildlife Service (Environment and Climate Change Canada), Ministry of Natural Resources and Forestry – Government of Ontario, Ontario Field Ornithologists (OFO), & Ontario Nature. (2009). *Ontario Breeding Birds Atlas*. <https://www.birdsontario.org/jsp/datasummaries.jsp>
- Bowfin Environmental Consulting Inc. (2021). *Cardinal Creek Village—South Side Headwater Drainage Feature Assessment*.
- California Academy of Sciences and National Geographic Society. (2024). *iNaturalist*. iNaturalist. <https://www.inaturalist.org/>
- City of Ottawa. (2021). *City of Ottawa Official Plan*. <https://ottawa.ca/en/planning-development-and-construction/official-plan-and-master-plans/official-plan#>
- City of Ottawa. (2022a). *Protocol for Wildlife Protection during Construction*. https://documents.ottawa.ca/sites/documents/files/documents/construction_en.pdf
- City of Ottawa. (2022b). *Significant Woodlands: Guidelines for Identification, Evaluation, and Impact Assessment*. https://documents.ottawa.ca/sites/documents/files/significant_woodlands_en.pdf
- City of Ottawa. (2023). *Environmental Impact Study Guidelines*. https://documents.ottawa.ca/sites/documents/files/eis_guidelines_tor_en.pdf
- City of Ottawa. (2024). *geoOttawa*. <https://maps.ottawa.ca/geoottawa/>
- City of Ottawa & AECOM. (2014). *Greater Cardinal Creek Subwatershed Management Plan (60189560)*. https://documents.ottawa.ca/sites/documents/files/documents/cardinal_creek_en.pdf
- COSEWIC. (2018). *COSEWIC assessment and status report on the Black Ash Fraxinus nigra in Canada* (p. xii + 95). Committee on the Status of Endangered Wildlife in Canada. <https://www.canada.ca/en/environment-climate-change/services/species-risk-public-registry/cosewic-assessments-status-reports/black-ash-2018.html>



DFO. (2023). *Aquatic Species at Risk Map*. Fisheries and Oceans Canada (Previously Department of Fisheries and Oceans, "DFO"). <https://www.dfo-mpo.gc.ca/species-especes/sara-lep/map-carte/index-eng.html>

Falk, B., Jokobsen, L., Surlykke, A., & Moss, C. F. (2014). *Bats coordinate sonar and flight behavior as they forage in open and cluttered environments*.
<https://journals.biologists.com/jeb/article/217/24/4356/12989/Bats-coordinate-sonar-and-flight-behavior-as-they>

Government of Canada. (1985). Fisheries Act, 1985 (R.S.C., 1985, c. F-14). <https://laws-lois.justice.gc.ca/eng/acts/f-14/>

Government of Canada. (1994). Migratory Birds Convention Act, 1994 (S.C. 1994, c. 22). <https://laws-lois.justice.gc.ca/eng/acts/m-7.01/>

Government of Canada. (2002). Species at Risk Act. 2002. S.C. 2002, c. 29.
<https://laws.justice.gc.ca/eng/acts/S-15.3/>

Government of Canada. (2024). *Species at Risk Public Registry*. <https://species-registry.canada.ca/index-en.html#/species?sortBy=commonNameSort&sortDirection=asc&pageSize=10>

Government of Ontario. (1990a). Conservation Authorities Act, R.S.O. 1990, c. C.27.
<https://www.ontario.ca/laws/statute/90c27>

Government of Ontario. (1990b). Planning Act, R.S.O. 1990, c. P.13.
<https://www.ontario.ca/laws/statute/90p13>

Government of Ontario. (1997). Fish and Wildlife Conservation Act, 1997, S.O. 1997, c. 41.
<https://www.ontario.ca/laws/statute/97f41>

Government of Ontario. (2007). Endangered Species Act. 2007. S.O. 2007, c.6.
<https://www.ontario.ca/laws/statute/07e06>



Government of Ontario. (2020). *Provincial Policy Statement, 2020—Under the Planning Act*.

<https://files.ontario.ca/mmah-provincial-policy-statement-2020-accessible-final-en-2020-02-14.pdf>

Humphrey, C. (2017). *Recovery Strategy for the Eastern Small-footed Myotis (Myotis leibii) in Ontario*

(Ontario Recovery Strategy Series, p. vii + 76). Prepared for the Ontario Ministry of Natural

Resources and Forestry. https://files.ontario.ca/mnrf_sar_rs_esfm_final_accessible.pdf

Humphrey, C., & Fotherby, H. (2019). *Recovery Strategy for the Little Brown Myotis (Myotis lucifugus),*

Northern Myotis (Myotis septentrionalis), and Tri-colored Bat (Perimyotis subflavus) in Ontario.

Adoption of the Recovery Strategy for the Little Brown Myotis (Myotis lucifugus), Northern Myotis

(Myotis septentrionalis), and Tri-colored Bat (Perimyotis subflavus) in Canada (Environment and

Climate Change Canada 2018). (Ontario Recovery Strategy Series, p. vii + 35). Prepared by the

Ministry of the Environment, Conservation and Parks. [https://files.ontario.ca/mecp-rs-bats-2019-](https://files.ontario.ca/mecp-rs-bats-2019-12-05.pdf)

[12-05.pdf](https://files.ontario.ca/mecp-rs-bats-2019-12-05.pdf)

Lee, H. R., Bakowsky, W., Riley, J., Bowles, J., Puddister, M., Uhlig, P., & McMurray, S. (1998). *Ecological*

Land Classification for Southern Ontario: First Approximation and its Application. Ontario Ministry

of Natural Resources. [https://www.researchgate.net/profile/Wasyl-](https://www.researchgate.net/profile/Wasyl-Bakowsky/publication/248626765_Ecological_Land_Classification_for_Southern_Ontario_First_Approximation_and_Its_Application/links/560e7abd08ae48337515fd59/Ecological-Land-Classification-for-Southern-Ontario-First-Approximation-and-Its-Application.pdf)

[Bakowsky/publication/248626765_Ecological_Land_Classification_for_Southern_Ontario_First_Approximation_and_Its_Application/links/560e7abd08ae48337515fd59/Ecological-Land-](https://www.researchgate.net/profile/Wasyl-Bakowsky/publication/248626765_Ecological_Land_Classification_for_Southern_Ontario_First_Approximation_and_Its_Application/links/560e7abd08ae48337515fd59/Ecological-Land-Classification-for-Southern-Ontario-First-Approximation-and-Its-Application.pdf)

[Classification-for-Southern-Ontario-First-Approximation-and-Its-Application.pdf](https://www.researchgate.net/profile/Wasyl-Bakowsky/publication/248626765_Ecological_Land_Classification_for_Southern_Ontario_First_Approximation_and_Its_Application/links/560e7abd08ae48337515fd59/Ecological-Land-Classification-for-Southern-Ontario-First-Approximation-and-Its-Application.pdf)

MECP. (2024). *Species at Risk in Ontario*. Ministry of the Environment, Conservation, and Parks.

<http://www.ontario.ca/page/species-risk-ontario>

MNR. (2010). *Natural Heritage Reference Manual for Natural Heritage Policies of the Provincial Policy*

Statement, 2005. Second Edition. Ministry of Natural Resources.

<https://docs.ontario.ca/documents/3270/natural-heritage-reference-manual-for-natural.pdf>



MNRF. (2015). *Significant Wildlife Habitat Criteria Schedules for Ecoregion 6E* (OMNRF Regional Operations Division: Southern Region Resources Section, p. 39). Ministry of Natural Resources and Forestry.

<https://dr6j45jk9xcmk.cloudfront.net/documents/4775/schedule-6e-jan-2015-access-ver-final-s.pdf>

MNRF. (2017). *Survey Protocol for Species at Risk Bats within Treed Habitats: Little Brown Myotis, Northern Myotis, and Tri-Coloured Bat* (p. 13). Ministry of Natural Resources and Forestry.

MNRF. (2024a). *Fish ON-Line*. Ministry of Natural Resources and Forestry.

<https://www.lioapplications.lrc.gov.on.ca/fishonline/Index.html?viewer=FishONLine.FishONLine&locale=en-CA>

MNRF. (2024b). *Land Information Ontario*. Ministry of Natural Resources and Forestry.

<http://www.ontario.ca/page/land-information-ontario>

MNRF. (2024c). *Natural Heritage Information Centre: Make a Natural Heritage Map*. Ministry of Natural Resources and Forestry. <http://www.ontario.ca/page/make-natural-heritage-area-map>

Ontario Breeding Bird Atlas. (2001). *Guide for Participants*. Atlas Management Board, Federation of Ontario Naturalists, Don Mills. https://www.birdsontario.org/jsp/download/obba_guide_en.pdf

Ontario Nature. (2019). *Ontario Reptile and Amphibian Atlas*.

<https://www.ontarioinsects.org/herp/index.html?Sort=0&area2=squaresCounties&records=all&myZoom=5&Lat=47.5&Long=-83.5>

Rideau Valley Conservation Authority. (2023). *Map A Property*. <https://www.rvca.ca/regulations-planning/map-a-property>

The Cornell Lab of Ornithology. (2024). *eBird: An online database of bird distribution and abundance*. <https://ebird.org/home>

Toronto and Region Conservation Authority & Credit Valley Conservation. (2013, July). *Evaluation, Classification and Management of Headwater Drainage Features Guidelines*. <https://trcaca.s3.ca->



central-1.amazonaws.com/app/uploads/2021/08/31112457/HDF-EVALUATION-CLASSIFICATION-
MANAGEMENT_2014.pdf

Toronto Entomologists' Association. (2024). *Ontario Butterfly Atlas*. <https://www.ontarioinsects.org/atlas/>
Wildlife Preservation Canada, The Xerces Society, The University of Ottawa, BeeSpotter, The Natural History
Museum, London, & the Montreal Insectarium. (2024). *Bumble Bee Watch: Bumble Sightings Map*.
https://www.bumblebeewatch.org/app/#/bees/map?filters=%7B%22sightingstatus_id%22:%5B%5D,%22species_id%22:%5B%2237%22%5D,%22province_id%22:%5B%5D%7D



Appendix A Qualifications of Report Authors



Maren Nielsen, BES, EMA

Maren is a Biologist with a background in terrestrial ecology. She has over eight years of comprehensive field, laboratory and environmental and agricultural consulting experience through a combination of graduate and undergraduate studies and work experience. Maren completed a Bachelor of Environmental Studies with Honours at York University and a Graduate Certificate in Environmental Management and Assessment from Niagara College Canada. Maren assists clients to navigate the land development and site rehabilitation processes as well as obtaining permits and approvals from regulatory agencies. She has led numerous studies including Environmental Assessments (EA), Environmental Impact Studies (EIS), Opportunities & Constraints Analysis, Agricultural Impact Assessments (AIA), LEAR Studies and Minimum Distance Separation (MDS) I & II studies. Maren has carried out field programs for the collection of soils, water, sediment, fish and benthos as well as vegetation surveys, wildlife surveys, wind turbine avian and bat mortality monitoring, and land use surveys. Since joining Kilgour & Associates Ltd. in 2023, Maren has worked on a variety of land development projects and completed numerous Environmental Impact Studies (EIS), Headwater Drainage Feature Assessments (HDFA), Existing Conditions Reports, Opportunities and Constraints Analysis, and Species at Risk (SAR) monitoring. Maren is a certified wetland evaluator under the Ontario Wetland Evaluation System (OWES).

Anthony Francis, PhD

Dr. Francis is a Senior Ecologist with 20 years' consulting experience to both government agencies and private industry. He has worked on a diversity of projects relating to species at risk (SAR), invasive species, terrestrial and aquatic habitat, environmental effects monitoring and mitigation, and fate/effects of contaminants. Within each of these subject areas, Dr. Francis has completed projects addressing specific site concerns and broader policy initiatives. Dr. Francis' academic background is in spatial ecology with a focus on tree species diversity. As a Senior Ecologist at KAL, he regularly completes TCRs, Environmental Impact Statements, and Integrated Environmental Reviews for land development projects throughout Ottawa and eastern Ontario. He is also a certified Butternut Health Assessor (BHA #104).



Appendix B Species at Risk Screening



Species Name (Taxonomic Name)	Status under Endangered Species Act (ESA)	Status under Schedule 1 of the Species at Risk Act (SARA)	Closest Species Occurrence Record to the Site	General Habitat Requirements	Site Suitability	Potential for Protected Elements ¹		Potential for Negative Interactions with Protected Elements ²
						Habitat	Individuals	
Birds								
Bald Eagle (<i>Haliaeetus leucocephalus</i>)	Special Concern	Not at Risk	Reported on-site (Cornell Lab of Ornithology, 2024)	Nest in mature forests near open water. In large trees such as pine and poplar.	Forested areas along the south tributary on-Site may provide suitable habitat.	Low	Low	Low
Bank Swallow (<i>Riparia riparia</i>)	Threatened	Threatened	Approximately 1.7 km from Site (Cornell Lab of Ornithology, 2024)	Colonial nester; burrows in eroding silt or sand banks, sand pit walls, and human-made sand piles. Often found on banks of rivers and lakes.	The fill piles and banks along the south tributary on-Site may provide suitable habitat.	Moderate	Moderate	Moderate
Barn Swallow (<i>Hirundo rustica</i>)	Special Concern	Threatened	Approximately 130 m from Site (Cornell Lab of Ornithology, 2024)	Nests on barns and other structures. Forages in open areas for flying insects. Lives in close association with humans and prefers to nest on structures such as open barns, under bridges, and in culverts.	Open areas on-Site may provide suitable foraging habitat. The Site does not contain suitable nesting habitat.	Negligible	Negligible	Negligible
Black Tern (<i>Chlidonias niger</i>)	Special Concern	Not at Risk	Approximately 1.7 km from Site (Cornell Lab of Ornithology, 2024)	Build floating nests in loose colonies in shallow marshes with abundant emergent vegetation, especially in cattails.	The Site does not appear to contain suitable habitat.	Negligible	Negligible	Negligible
Bobolink (<i>Dolichonyx oryzivorus</i>)	Threatened	Threatened	Approximately 1.1 km from Site (Cornell Lab of Ornithology, 2024)	Breeds in hayfields, pastures, agricultural fields, and abandoned fields with tall grass that are ≥5 ha, and preferably >30 ha.	Agricultural fields on-Site may provide suitable habitat.	Moderate	Moderate	Moderate
Canada Warbler (<i>Cardellina canadensis</i>)	Special Concern	Threatened	Approximately 400 m from Site (Cornell Lab of Ornithology, 2024)	Prefers moist forests with dense shrub layers. Nests located on or near the ground on mossy logs or roots, along stream banks or on hummocks. Area-sensitive species that usually require a minimum of 30 ha of continuous forest for breeding habitat (OMNR, 2000).	Forested areas along the south tributary on-Site may provide suitable habitat.	Moderate	Moderate	Moderate
Chimney Swift (<i>Chaetura pelagica</i>)	Threatened	Threatened	Approximately 4 km from site (Cornell Lab of Ornithology, 2024)	Nests in traditional-style open brick chimneys (and rarely in hollow trees). Tends to stay close to water.	The Site does not appear to contain suitable habitat.	Negligible	Negligible	Negligible



Species Name (Taxonomic Name)	Status under Endangered Species Act (ESA)	Status under Schedule 1 of the Species at Risk Act (SARA)	Closest Species Occurrence Record to the Site	General Habitat Requirements	Site Suitability	Potential for Protected Elements ¹		Potential for Negative Interactions with Protected Elements ²
						Habitat	Individuals	
Common Nighthawk (<i>Chordeiles minor</i>)	Special Concern	Threatened	Approximately 2 km from Site (Cornell Lab of Ornithology, 2024)	Nests in a wide variety of open sites, including beaches, fields, and gravel rooftops with little to no ground vegetation. They also nest in cultivated fields, orchards, urban parks, mine tailings and along gravel roads/railways but tend to occupy more natural sites.	Open, agricultural areas on-Site may provide suitable habitat.	Low	Low	Low
Eastern Meadowlark (<i>Sturnella magna</i>)	Threatened	Threatened	Approximately 130 m from Site (Cornell Lab of Ornithology, 2024)	Breeds in hayfields, pastures, agricultural fields, and abandoned fields with tall grass that are ≥5 ha, and preferably >30 ha.	Agricultural fields on-Site may provide suitable habitat.	Moderate	Moderate	Moderate
Eastern Whip-poor-will (<i>Antrostomus vociferus</i>)	Threatened	Threatened	Approximately 1.7 km from Site (Cornell Lab of Ornithology, 2024)	Suitable breeding habitats generally include open and half treed areas and often exhibit a scattered distribution of treed and open space. Lays eggs directly on the forest floor. Roosts are typically located in forest habitat on a low branch or directly on the ground. Home range size varies from 20 to 500 ha (mean 136 ha) (ECCC, 2018a).	Open areas and forested patches along the south tributary on-Site may provide suitable habitat	Low	Low	Low
Eastern Wood-Pewee (<i>Contopus virens</i>)	Special Concern	Special Concern	Approximately 1.5 km from Site (Cornell Lab of Ornithology, 2024)	Woodland species often found in the mid-canopy layer near clearings and edges of intermediate age and mature deciduous and mixed forests with little understory.	Forested areas along the south tributary on-Site may provide suitable habitat.	Moderate	Moderate	High Detected onsite
Evening Grosbeak (<i>Coccothraustes vespertinus</i>)	Special Concern	Special Concern	Approximately 1.8 km from Site (Cornell Lab of Ornithology, 2024)	Nests in trees or large shrubs. Prefers mature coniferous forests (fir and/or spruce dominated), but will also use deciduous forests, parklands, and orchards. Its abundance is strongly linked to the cycle of Spruce Budworm.	Forested areas along the south tributary on-Site may provide suitable habitat.	Low	Low	Low
Golden Eagle (<i>Aquila chrysaetos</i>)	Endangered	Not at Risk	Approximately 1.7 km from Site (Cornell Lab of Ornithology, 2024)	Nests in remote, undisturbed areas, usually building their nests on ledges on a steep cliff/riverbank or large trees if	Steep riverbanks along the south tributary may provide suitable habitat.	Low	Low	Low



Species Name (Taxonomic Name)	Status under Endangered Species Act (ESA)	Status under Schedule 1 of the Species at Risk Act (SARA)	Closest Species Occurrence Record to the Site	General Habitat Requirements	Site Suitability	Potential for Protected Elements ¹		Potential for Negative Interactions with Protected Elements ²
						Habitat	Individuals	
				needed. Most hunting is done near open areas such as large bogs or tundra. Migration only; no reported nests in Ottawa.				
Hudsonian Godwit (<i>Limosa haemastica</i>)	Threatened	No Status	Approximately 1.7 km from Site (Cornell Lab of Ornithology, 2024)	They use a wide variety of habitats during migration, such as freshwater marshes, saline lakes, flooded fields, shallow ponds, coastal wetlands, and mudflats. Migrant only; breeds in far north.	The Site does not appear to contain suitable habitat.	Negligible	Negligible	Negligible
Least Bittern (<i>Ixobrychus exilis</i>)	Threatened	Threatened	Approximately 1.7 km from Site (Cornell Lab of Ornithology, 2024)	Found in a variety of wetland habitats, but strongly prefers cattail marshes with a mix of open pools and channels. They prefer larger marshes >5 ha in size and are intolerant of loss of habitat and human disturbance (OMNR, 2000).	The Site does not appear to contain suitable habitat.	Negligible	Negligible	Negligible
Lesser Yellowlegs (<i>Tringa flavipes</i>)	Threatened	No Status	Approximately 1.8 km from Site (Cornell Lab of Ornithology, 2024)	Breeds in boreal wetlands. Nests on dry ground or forest openings near peatlands, marshes, and ponds in the boreal forest and taiga (Government of Canada, 2021). Migrant only; nests in far north.	The Site does not appear to contain suitable habitat.	Negligible	Negligible	Negligible
Olive-sided Flycatcher (<i>Contopus cooperi</i>)	Special Concern	Threatened	Approximately 1.7 km from Site (Cornell Lab of Ornithology, 2024)	Found along coniferous or mixed forest edges and openings. Will use forests that have been logged or burned if there are ample tall snags and trees to use for foraging perches.	The forested areas along the south tributary on-Site may contain suitable habitat.	Moderate	Moderate	Moderate
Peregrine Falcon (<i>Falco peregrinus</i>)	Special Concern	Special Concern	Approximately 2 km from Site (Cornell Lab of Ornithology, 2024)	Nests on tall, steep cliff ledges close to large bodies of water. Urban peregrines raise their young on ledges of tall buildings, even in busy downtown areas.	The Site does not appear to contain suitable habitat.	Negligible	Negligible	Negligible
Red Knot (<i>Calidris canutus rufa</i>)	Endangered	Endangered	Approximately 1.8 km from Site (Cornell Lab of Ornithology, 2024)	Prefer open beaches, mudflats, and coastal lagoons where they feast on molluscs, crustaceans, and other invertebrates. Migrant only; nests in far north.	The Site does not appear to contain suitable habitat.	None	None	None
Rusty Blackbird (<i>Euphagus carolinus</i>)	Special Concern	Special Concern	Approximately 1.7 km from Site (Cornell Lab of Ornithology, 2024)	Prefers wet wooded or shrubby areas. Nests at edges of boreal wetlands and coniferous forests. These areas include bogs, marshes, and beaver ponds.	The Site does not appear to contain suitable habitat.	Negligible	Negligible	Negligible



Species Name (Taxonomic Name)	Status under Endangered Species Act (ESA)	Status under Schedule 1 of the Species at Risk Act (SARA)	Closest Species Occurrence Record to the Site	General Habitat Requirements	Site Suitability	Potential for Protected Elements ¹		Potential for Negative Interactions with Protected Elements ²
						Habitat	Individuals	
Short-eared Owl (<i>Asio flammeus</i>)	Threatened	Special Concern	Within 5 km of Site (MNR, 2024a)	Prefer a mosaic of grasslands and wetlands. Lives in open areas such as grasslands, marshes, and tundra where it nests on the ground and hunts for small mammals (Environment Canada, 2016c).	The Site does not appear to contain suitable habitat.	Negligible	Negligible	Negligible
Wood Thrush (<i>Hylocichla mustelina</i>)	Special Concern	Threatened	Within 10 km of Site (Birds Canada et al., 2009)	Lives in mature deciduous and mixed forests. They seek moist stands of trees with well-developed undergrowth and tall trees for singing and perching. Prefers nesting in large forest mosaics, but will also use fragmented forests. Usually build nests in Sugar Maple or American Beech.	Forested areas along the south tributary on-Site may provide suitable habitat.	Moderate	Moderate	Moderate
Mammals								
Eastern Small-footed Myotis (<i>Myotis leibii</i>)	Endangered	Not Listed	Humphrey (2017) – in region	In the spring and summer, Eastern Small-footed Myotis will roost in a variety of habitats, including in or under rocks, in rock outcrops, in buildings, under bridges, or in caves, mines, or hollow trees. Overwinters in caves and abandoned mines.	Forested areas along the south tributary on-Site may provide suitable roosting habitat.	Moderate	Moderate	Moderate
Little Brown Myotis (<i>Myotis lucifugus</i>)	Endangered	Endangered	Humphrey and Fotherby (2019) – in region	During the day they roost in trees and buildings. They often select attics, abandoned buildings, and barns for summer colonies where they can raise their young. They can squeeze through very tiny spaces (as small as six millimetres across) allowing them access to many different roosting areas.	Forested areas along the south tributary on-Site may provide suitable roosting habitat.	Moderate	Moderate	Moderate
Northern Myotis / Northern Long-eared Bat (<i>Myotis septentrionalis</i>)	Endangered	Endangered	Humphrey and Fotherby (2019) – in region	Associated with deciduous and mixed forests, choosing to roost under loose bark and in the cavities of trees. They forage along and within forests as well as in hayfields and pastures adjacent to mixed forests.	Forested areas along the south tributary on-Site may provide suitable roosting habitat.	Moderate	Moderate	Moderate



Species Name (Taxonomic Name)	Status under Endangered Species Act (ESA)	Status under Schedule 1 of the Species at Risk Act (SARA)	Closest Species Occurrence Record to the Site	General Habitat Requirements	Site Suitability	Potential for Protected Elements ¹		Potential for Negative Interactions with Protected Elements ²
						Habitat	Individuals	
Tri-colored Bat / Eastern Pipistrelle (<i>Perimyotis subflavus</i>)	Endangered	Endangered	Humphrey and Fotherby (2019) – in region	Roosts mainly in trees during summer; overwinters in caves and mines along with other species, but often uses deeper parts of the hibernaculum. Foraging occurs in forested riparian areas, over water, and within gaps in forest canopies.	Forested areas along the south tributary on-Site may provide suitable roosting habitat.	Moderate	Moderate	Moderate
Hoary Bat (<i>Lasiurus cinereus</i>)	Endangered (January 2025)	No Status	n/a	Roosts in both deciduous and coniferous forests of any age, among canopy foliage with open flight space below. Maternity roosts are often in large diameter, tall trees. Foraging occurs in open areas, wetlands, grasslands and open fields, with sparse trees.	Habitat on site is generally suitable.	Moderate	Moderate	Moderate
Silver-haired Bat (<i>Lasionycteris noctivagans</i>)	Endangered (January 2025)	No Status	n/a	Roosts under bark and in large decaying deciduous and coniferous tree cavities. Foraging occurs in young and mature forest openings and along forest edges.	Habitat on site is generally suitable.	Moderate	Moderate	Moderate
Eastern Red Bat (<i>Lariurus borealis</i>)	Endangered (January 2025)	No Status	n/a	Roosts in both deciduous and coniferous forests of any age, among canopy foliage with open flight space below. Maternity roosts are often in large diameter, tall trees. Foraging occurs in forested and non-forested areas, above and below forest canopies.	Habitat on site is generally suitable.	Moderate	Moderate	Moderate
Amphibians								
Western Chorus Frog (<i>Pseudacris triseriata</i>)	Not Listed	Great Lakes/ St. Lawrence population: Threatened	Within 5 km of Site (MNR, 2024a)	Inhabits forest openings around woodland ponds but can also be found in or near damp meadows, marshes, bottomland swamps, and temporary ponds in open country, or even urban areas.	Wetland area near the south edge of the Site and associated forested areas may provide suitable habitat.	Low	Low	Low
Reptiles								
Blanding's Turtle (<i>Emydoidea blandingii</i>)	Threatened	Endangered	Within 5 km of Site (MNR, 2024a)	Quiet lakes, streams, and wetlands with abundant emergent vegetation. Also frequently occurs in adjacent upland forests.	The Site does not appear to contain suitable habitat.	Low	Low	Low
Eastern Milksnake	Not Listed	Special Concern	Within 10 km of Site (Ontario Nature, 2019)	Found in a variety of open and edge habitats, including	Forest edges and open areas on-Site may provide suitable habitat.	Low	Low	Low



Species Name (Taxonomic Name)	Status under Endangered Species Act (ESA)	Status under Schedule 1 of the Species at Risk Act (SARA)	Closest Species Occurrence Record to the Site	General Habitat Requirements	Site Suitability	Potential for Protected Elements ¹		Potential for Negative Interactions with Protected Elements ²
						Habitat	Individuals	
<i>Lampropeltis triangulum</i>				meadows, rocky outcrops, and forest edges. They can also inhabit forests. Further, they are often associated with human-made structures such as barns (Environment Canada, 2015b).				
Eastern Musk Turtle / Stinkpot (<i>Sternotherus odoratus</i>)	Special Concern	Special Concern	Within 10 km of Site (Ontario Nature, 2019)	Found in lakes, ponds, marshes, and rivers that are generally slow-moving, have abundant emergent vegetation, and muddy bottoms that they burrow into for winter hibernation.	The Site does not appear to contain suitable habitat.	Low	Low	Low
Midland Painted Turtle (<i>Chrysemys picta marginata</i>)	Not Listed	Special Concern	Within 5 km of Site (MNR, 2024a)	Inhabits waterbodies, such as ponds, marshes, lakes, and slow-moving creeks that have a soft bottom and provide abundant basking sites and aquatic vegetation. Often bask on shorelines or on logs and rocks that protrude from the water.	The south tributary on-Site may provide suitable habitat.	Low	Low	Low
Northern Map Turtle (<i>Graptemys geographica</i>)	Special Concern	Special Concern	Within 5 km of Site (MNR, 2024a)	Lives in rivers and lakeshores where it basks on emergent rocks and fallen trees throughout the spring and summer. In winter, they hibernate on the bottom of deep, slow-moving sections of river.	The Site does not appear to contain suitable habitat.	Low	Low	Low
Snapping Turtle (<i>Chelydra serpentina</i>)	Special Concern	Special Concern	Within 5 km of Site (MNR, 2024a)	Spend most of their lives in the water. Prefer shallow waters so they can hide under the soft mud and leaf litter with only their noses exposed to the surface to breathe.	The Site does not appear to contain suitable habitat.	Low	Low	Low
Arthropods								
Monarch (<i>Danaus plexippus</i>)	Special Concern	Special Concern	Within 10 km of Site (Toronto Entomologists' Association, 2024)	Milkweeds are the sole food plant for Monarch caterpillars. These plants predominantly grow in open and periodically disturbed habitats such as roadsides, fields, wetlands, prairies, and open forests.	Open meadows and forest edges may provide suitable habitat.	Negligible	Negligible	Negligible



Species Name (Taxonomic Name)	Status under Endangered Species Act (ESA)	Status under Schedule 1 of the Species at Risk Act (SARA)	Closest Species Occurrence Record to the Site	General Habitat Requirements	Site Suitability	Potential for Protected Elements ¹		Potential for Negative Interactions with Protected Elements ²
						Habitat	Individuals	
Yellow-banded Bumble Bee (<i>Bombus terricola</i>)	Special Concern	Special Concern	Within 5 km of Site (MNRF, 2024a)	This species is a forage and habitat generalist, able to use a variety of nectaring plants and environmental conditions. Can be found in mixed woodlands, particularly for nesting and overwintering, as well as a variety of open habitat such as native grasslands, farmlands, and urban areas.	Forested areas along the south tributary on-Site may provide suitable habitat.	Low	Low	Low
Vascular Plants								
Black Ash (<i>Fraxinus nigra</i>)	Endangered	No Status	Onsite (KAL, 2024)	Predominantly a wetland species found in swamps, floodplains, and fens.	The Site does not appear to contain suitable habitat.	High	High	High
Butternut (<i>Juglans cinerea</i>)	Endangered	Endangered	Within 5 km of Site (MNRF, 2024a)	Commonly found in riparian habitats but is also found on rich, moist, well-drained loams and well-drained gravels, especially those of limestone origin.	Riparian forests along the south tributary may provide suitable habitat.	Moderate	Moderate	Low Not observed on the Site

1 The potential for occurrence of protected habitats and individuals within the project area is estimated based on the following considerations:

	Habitat	Individuals
None	It is not possible for the habitat of the species to occur in proximity to the project site	The species is documented as no longer occurring in the ecoregion or could not occur in proximity to the project area.
Negligible	The usage of the project site as habitat is possible but would be highly unlikely/unusual.	Transient occurrence near the project area is possible but is very unlikely.
Low	The project site includes areas that could be used by the species as habitat, but such usage is considered unlikely given the quality of the feature, a lack of individuals in the broader area, or other (relative) site considerations.	Transient occurrence near the project area possible, but the species would be unlikely to use or require the area.
Moderate	The project site includes areas that could reasonably be expected to provide confirmed or defined habitat within a time frame relevant to the project.	The species occurs in the vicinity and could actively use the site, or transient occurrence should be anticipated.
High	The project site includes areas confirmed to actively provide habitat or to constitute habitat based on official habitat description guidance documents.	The species is confirmed as present on, and actively using the site.

2 The potential for negative project interaction with species and/or their habitat is estimated considering both the likelihood of presence and the general details of the project (e.g., timing, extent), and following the definitions below. If the potential differs for habitat and individuals, the higher value is reported, unless otherwise justified

	Habitat	Individuals
None	It is not possible for the species to occupy the site area due to access barriers.	The species is documented as no longer occurring in the ecoregion
Negligible	Negligible habitat potential, or low habitat potential and the project would not be anticipated to alter the habitat.	Negligible occurrence potential for presence, or absence during the entire span of the project.
Low	Low habitat potential, or medium habitat potential and the project would not be anticipated to alter the habitat.	Low occurrence potential for presence, or the project design excludes individuals in a non-harassing manner by default.



Moderate	Medium habitat potential, or high habitat potential and the project would not be anticipated to alter the habitat (as expressed by MECF).	Medium occurrence potential for presence, or the project design excludes individuals in accordance with agency guidelines/directives by default (i.e., outside of mitigation measures prescribed in this report).
High	The project area will alter identified habitat.	The project will interact with individuals.



LITERATURE CITED

- Birds Canada, Canadian Wildlife Service (Environment and Climate Change Canada), Ministry of Natural Resources and Forestry – Government of Ontario, Ontario Field Ornithologists (OFO), and Ontario Nature. 2009. Atlas of the Breeding Birds of Ontario 2001-2005. Available online at: <https://www.birdsontario.org/jsp/datasummaries.jsp>
- Burke, Peter S. 2012. Management Plan for the Black Tern (*Chlidonias niger*) in Ontario. Ontario Management Plan Series. Prepared for the Ontario Ministry of Natural Resources (OMNR), Peterborough, Ontario. vi + 47 pp. Available online at: https://files.ontario.ca/environment-and-energy/species-at-risk/mnr_sar_mtpln_blktrn_en.pdf
- California Academy of Sciences and National Geographic Society. 2022. iNaturalist. Available online at: <https://www.inaturalist.org/>
- Committee on the Status of Endangered Wildlife in Canada. 2011. COSEWIC Assessment and Status Report on the Macropis Cuckoo Bee (*Epeoloides pilosulus*) in Canada. Available online at: https://www.registrelep-sararegistry.gc.ca/virtual_sara/files/cosewic/sr_macropis_cuckoo_bee_0911_eng.pdf
- Committee on the Status of Endangered Wildlife in Canada. 2015. COSEWIC Assessment and Status Report on the Black-foam Lichen (*Anzia colpodes*) in Canada. Available online at: https://sararegistry.gc.ca/virtual_sara/files/cosewic/sr_Black-foam%20Lichen_2015_e.pdf
- The Cornell Lab of Ornithology. 2022. eBird: An online database of bird distribution and abundance. Available online at: <https://ebird.org/home>
- COSEWIC. 2018. COSEWIC assessment and status report on the American Bumble Bee *Bombus pensylvanicus* in Canada. Committee on the Status of Endangered Wildlife in Canada. Ottawa. x + 52 pp. Available online at: <https://www.canada.ca/en/environment-climate-change/services/species-risk-public-registry/cosewic-assessments-status-reports/american-bumble-bee-2018.html>
- COSEWIC. 2019. COSEWIC assessment and status report on the Suckley's Cuckoo Bumble Bee *Bombus suckleyi* in Canada. Committee on the Status of Endangered Wildlife in Canada. Ottawa. xi + 70 pp. Available online at: <https://www.canada.ca/en/environment-climate-change/services/species-risk-public-registry/cosewic-assessments-status-reports/suckley-cuckoo-bumble-bee-2019.html>
- Environment Canada. 2011. Management Plan for the Louisiana Waterthrush (*Seiurus motacilla*) in Canada [Proposed]. Species at Risk Act Management Plan Series. Environment Canada, Ottawa. iii + 18 pp. Available online at: https://www.sararegistry.gc.ca/virtual_sara/files/plans/mp_louisiana_waterthrush_e.pdf
- Environment Canada. 2013. Management Plan for the Yellow Rail (*Coturnicops noveboracensis*) in Canada. Species at Risk Act Management Plan Series. Environment Canada, Ottawa. iii + 24 pp. Available online at: https://www.sararegistry.gc.ca/virtual_sara/files/plans/mp_yellow_rail_e_final.pdf
- Environment Canada. 2014a. Recovery Strategy for the Western Chorus Frog (*Pseudacris triseriata*), Great Lakes / St. Lawrence – Canadian Shield Population, in Canada [Proposed], Species at Risk Act Recovery Strategy Series, Environment Canada, Ottawa, v + 46 pp. Available online at: https://www.registrelep-sararegistry.gc.ca/virtual_sara/files/plans/rs_rainette-fx-grillon-ouest-w-chorus-frog-prop-0614_e.pdf



- Environment Canada. 2014b. Management Plan for the Eastern Ribbonsnake (*Thamnophis sauritus*), Great Lakes population, in Canada [Proposed]. Species at Risk Act Management Plan Series. Environment Canada, Ottawa. iv + 23 pp. Available online at: https://www.sararegistry.gc.ca/virtual_sara/files/plans/mp_eastern_ribbonsnake_e_proposed.pdf
- Environment Canada. 2015a. Management Plan for the Rusty Blackbird (*Euphagus carolinus*) in Canada. Species at Risk Act Management Plan Series. Environment Canada, Ottawa. iv + 26 pp. Available online at: https://www.sararegistry.gc.ca/virtual_sara/files/plans/mp_rusty_blackbird_e_final.pdf
- Environment Canada. 2015b. Management Plan for the Eastern Milksnake (*Lampropeltis triangulum*) in Canada. Species at Risk Act Management Plan Series. Environment Canada, Ottawa. iii + 27 pp. Available online at: https://www.canada.ca/en/environment-climate-change/services/species-risk-public-registry/management-plans/eastern-milksnake-2015.html#_03_3
- Environment Canada. 2016a. Recovery Strategy for the Canada Warbler (*Cardellina canadensis*) in Canada. Species at Risk Act Recovery Strategy Series. Environment Canada, Ottawa. vii + 56 pp. Available online at: https://www.sararegistry.gc.ca/virtual_sara/files/plans/rs_canada%20warbler_e_final.pdf
- Environment Canada. 2016b. Recovery Strategy for the Common Nighthawk (*Chordeiles minor*) in Canada. Species at Risk Act Recovery Strategy Series. Environment Canada, Ottawa. vii + 49 pp. Available online at: https://www.registrelep-sararegistry.gc.ca/virtual_sara/files/plans/rs_common%20nighthawk_e_final.pdf
- Environment Canada. 2016c. Management Plan for the Short-eared Owl (*Asio flammeus*) in Canada [Proposed]. Species at Risk Act Management Plan Series. Environment Canada, Ottawa. v + 35 pp. Available online at: file:///C:/Users/skatsaras/Downloads/Management_plan_asio_falmmuus_Feb23_2016.pdf
- Environment and Climate Change Canada (ECCC). 2018a. Recovery Strategy for the Eastern Whip-poor-will (*Antrastomus vociferus*) in Canada. Species at Risk Act Recovery Strategy Series. Environment and Climate Change Canada, Ottawa. vi + 107 pp. Available online at: <https://files.ontario.ca/mecp-rs-easternwhip-poor-will-2019-12-05.pdf>
- Environment and Climate Change Canada (ECCC). 2018b. Recovery Strategy for the Blanding's Turtle (*Emydoidea blandingii*), Great Lakes / St. Lawrence population, in Canada. Species at Risk Act Recovery Strategy Series. Environment and Climate Change Canada, Ottawa. viii + 59 pp. Available online at: <https://files.ontario.ca/mecp-rs-blandings-turtle-2019-12-05.pdf>
- Environment and Climate Change Canada (ECCC). 2022. Management Plan for the Yellow banded Bumble Bee (*Bombus terricola*) in Canada [Proposed]. Species at Risk Act Management Plan Series. Environment and Climate Change Canada, Ottawa. iv + 46 pp. Available online at: <https://www.canada.ca/en/environment-climate-change/services/species-risk-public-registry/management-plans/yellow-banded-bumble-bee-proposed-2022.html>
- Fisheries and Oceans Canada (previously Department of Fisheries and Oceans, "DFO"). 2022. Aquatic Species at Risk Map. Available online at: <https://www.dfo-mpo.gc.ca/species-especes/sara-lep/map-carte/index-eng.html>
- Government of Canada. 2019. American Bumble Bee (*Bombus pensylvanicus*): COSEWIC assessment and status report 2018. Available online at: <https://www.canada.ca/en/environment-climate-change/services/species-risk-public-registry/cosewic-assessments-status-reports/american-bumble-bee-2018.html>
- Government of Canada. 2020. Suckley's Cuckoo Bumble Bee (*Bombus suckleyi*): COSEWIC assessment and status report 2019. Available online at: <https://www.canada.ca/en/environment-climate->



[change/services/species-risk-public-registry/cosewic-assessments-status-reports/suckley-cuckoo-bumble-bee-2019.html](https://www2.gov.bc.ca/gov2/serv/bc/change/services/species-risk-public-registry/cosewic-assessments-status-reports/suckley-cuckoo-bumble-bee-2019.html)

Government of Canada. 2021. Lesser Yellowlegs (*Tringa flavipes*): COSEWIC assessment and status report 2020. Available online at: <https://www.canada.ca/en/environment-climate-change/services/species-risk-public-registry/cosewic-assessments-status-reports/lesser-yellowlegs-2020.html>

Government of Canada. 2022. Species at Risk Public Registry. Available online at: http://www.registrelep-sararegistry.gc.ca/sar/index/default_e.cfm

Humphrey, C. 2017. Recovery Strategy for the Eastern Small-footed Myotis (*Myotis leibii*) in Ontario. Ontario Recovery Strategy Series. Prepared for the Ontario Ministry of Natural Resources and Forestry, Peterborough, Ontario. vii + 76 pp. Available online at: https://files.ontario.ca/mnrf_sar_rs_esfm_final_accessible.pdf

Humphrey, C., and H. Fotherby. 2019. Recovery Strategy for the Little Brown Myotis (*Myotis lucifugus*), Northern Myotis (*Myotis septentrionalis*) and Tri-colored Bat (*Perimyotis subflavus*) in Ontario. Ontario Recovery Strategy Series. Prepared by the Ministry of the Environment, Conservation and Parks, Peterborough, Ontario. vii + 35 pp. + Appendix. Adoption of the Recovery Strategy for the Little Brown Myotis (*Myotis lucifugus*), the Northern Myotis (*Myotis septentrionalis*), and the Tri-colored Bat (*Perimyotis subflavus*) in Canada (Environment and Climate Change Canada 2018). Available online at: <https://files.ontario.ca/mecp-rs-bats-2019-12-05.pdf>

Kraus, T., B. Hutchinson, S. Thompson and K. Prior. 2010. Recovery Strategy for the Gray Ratsnake (*Pantherophis spiloides*) – Carolinian and Frontenac Axis populations in Ontario. Ontario Recovery Strategy Series. Prepared for the Ontario Ministry of Natural Resources, Peterborough, Ontario. vi + 23 pp. Available online at: <https://www.ontario.ca/page/gray-ratsnake-recovery-strategy>

Ministry of Environment, Conservation, and Parks (MECP). 2019a. Algonquin Wolf. Available online at: <https://www.ontario.ca/page/algonquin-wolf>

Ministry of Environment, Conservation, and Parks (MECP). 2019b. Gypsy Cuckoo Bumble Bee. Available online at: <https://www.ontario.ca/page/gypsy-cuckoo-bumble-bee>

Ministry of Environment, Conservation, and Parks (MECP). 2019c. Nine-spotted Lady Beetle. Available online at: <https://www.ontario.ca/page/nine-spotted-lady-beetle>

Ministry of Environment, Conservation, and Parks (MECP). 2019d. Rapids Clubtail. Available online at: <https://www.ontario.ca/page/rapids-clubtail>

Ministry of Environment, Conservation, and Parks (MECP). 2019e. Rusty-patched Bumble Bee. Available online at: <https://www.ontario.ca/page/rusty-patched-bumble-bee>

Ministry of Environment, Conservation, and Parks (MECP). 2019f. Pale-bellied Frost Lichen. Available online at: <https://www.ontario.ca/page/pale-bellied-frost-lichen>

Ministry of Environment, Conservation, and Parks (MECP). 2020a. Gray Fox. Available online at: <https://www.ontario.ca/page/grey-fox>

Ministry of Environment, Conservation, and Parks (MECP). 2020b. Transverse Lady Beetle. Available online at: <https://www.ontario.ca/page/transverse-lady-beetle>



Ministry of Environment, Conservation, and Parks (MECP). 2022. Species at Risk in Ontario. Available online at:
<https://www.ontario.ca/page/species-risk-ontario>

Ministry of Natural Resources and Forestry (MNRF). 2022a. Natural Heritage Information Centre: Make Natural Heritage Map. Available online at: <https://www.ontario.ca/page/make-natural-heritage-area-map>

Ministry of Natural Resources and Forestry (MNRF). 2022b. Land Information Ontario. Available online at:
<https://www.ontario.ca/page/land-information-ontario>

Ontario Ministry of Natural Resources (OMNR). 2000. Significant Wildlife Habitat Technical Guide. Fish and Wildlife Branch. Wildlife Section. Science Development and Transfer Branch. Southcentral Sciences Section. 151 pp. Available at: <https://dr6j45jk9xcmk.cloudfront.net/documents/3620/significant-wildlife-habitat-technical-guide.pdf>

Ontario Nature. 2019. Ontario Reptile and Amphibian Atlas. Available online at:
<https://www.ontarioinsects.org/herp/index.html?Sort=0&area2=squaresCounties&records=all&myZoom=5&Lat=47.5&Long=-83.5>

Seburn, D.C. 2010. Recovery strategy for the Common Five-lined Skink (*Plestiodon fasciatus*) – Carolinian and Southern Shield populations in Ontario. Ontario Recovery Strategy Series. Prepared for the Ontario Ministry of Natural Resources, Peterborough, Ontario. vi + 22 pp. Available online at:
<https://www.ontario.ca/page/common-five-lined-skink-recovery-strategy>

Toronto Entomologists' Association. 2022. Ontario Butterfly Atlas. Available online at:
<https://www.ontarioinsects.org/atlas/>

Wildlife Preservation Canada, the Xerces Society, the University of Ottawa, BeeSpotter, The Natural History Museum, London, and the Montreal Insectarium. 2022. Bumble Bee Watch: Bumble Sightings Map. Available online at:
https://www.bumblebeewatch.org/app/#/bees/map?filters=%7B%22sightingstatus_id%22:%5B%5D,%22species_id%22:%5B%2237%22%5D,%22province_id%22:%5B%5D%7D

